

(10) International Publication Number  
WO 02/04520 A2

Roderick, T. [TUGB]: 8 Boundary Court, St. Lawrence

Jayolaxmi JIN/USI: 34,359 Mayhird Citele, Fremont, CA  
94555 (US), 141. Yan ICN/USI: 3885 Corona Way, Palmdale, CA 93550 (US)

233 Coy Drive, San Jose, CA 95123 (US). AZIMZAI

Valley, CA 94552 (U.S.). I.A.L., Preeti [IN/US]; P.O. Box

USUSJ: 3770 Polton Place Way, San Jose, CA 95121

Drive, San Jose, CA 95118 (US). XL, Yumling JUS/USJ  
1720 Walnut Drive Mountain View CA 94039 (US)

SELHAMER, Jeffrey, J. (US/US): 12535 La Cresta  
1st Alvar Hills, CA 94027 (US) BOBOWSKY, Mark

L | US/USI: 122 Orchard Avenue, Redwood City, CA 94061 (US). KHAN, Farrah. A. | IN/USI: 3617 Centura

11/US/US): 50 Woodside Avenue, San Francisco, CA 94127

locio Avenue #23, Mountain View, CA 94043 (US). DAS

Mountain View, CA 94040 (US). POLICKY, Jennifer, L.

# 11001 EST COV Diets and Food Consumption

Inc., 3160 Porter Drive, Palo Alto, CA 94304 (U.S.).

Designated States (national): AE, AG, AL, AM, AT, AU, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, FR, GB, GR, GU, HK, HN, HU, IL, IN, JP, KE, KG, KH, KR, KZ, LA, LB, LC, LI, LU, LV, MA, MG, MK, MN, MU, MV, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, QA, RO, RU, RW, SA, SC, SD, SE, SG, SI, SK, SL, SM, SN, SR, ST, SV, SY, TD, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VE, VN, YU, ZA, ZM, ZW.

CA, CH, CL, CM, CN, CO, CR, CU, CY, CZ, DE, DK, DM, DO, EC, EE, EG, ES, FI, FR, GB, GR, GT, HK, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LI, LU, LV, LY, MA, MC, MD, ME, MG, MK, MN, MU, MV, MW, MY, MZ, NA, NE, NG, NI, NL, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, QA, RO, RU, RW, SA, SB, SC, SD, SE, SG, SI, SK, SL, SN, SR, ST, SV, TD, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UY, UZ, VC, VE, VN, VU, WF, YU, ZA, ZM, ZW.

MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL

100

KE, IS, MW, MZ, SD, SI, SZ, TZ, UC, ZW), Eurasia

patient (AT, BL, CL, CY, DL, DK, LS, FL, IR, GB, GR, IT, LL, MC, NL, PT, SF, TR). OAP patient (BE, FI, CL)

CG, CI, CM, GA, GN, GW, ML, MK, NE, SN, ID, IG).

*without international search report and to be republished*

up on receipts of mail: open:

Notes on Codes and Abbreviations" appearing at the begin-

---

ills, antibodies, agonists, and antagonists. The invention also

## TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of transporters and ion channels and to the use of these sequences in the diagnosis, treatment, and prevention of transport, neurological, muscle, immunological, and cell proliferative disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of transporters and ion channels.

## BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded and subdivided into functionally distinct organelles by hydrophobic lipid bilayer membranes which are highly impermeable to most polar molecules. Cells and organelles require transport proteins to import and export essential nutrients and metal ions including  $\text{K}^+$ ,  $\text{NH}_4^+$ ,  $\text{P}_i$ ,  $\text{SO}_4^{2-}$ , sugars, and vitamins, as well as various metabolic waste products. Transport

proteins also play roles in antibiotic resistance, toxin secretion, ion balance, synaptic neurotransmission, kidney function, intestinal absorption, tumor growth, and other diverse cell functions (Griffith, J. and C. Sausom (1998) The Transporter Facts Book, Academic Press, San Diego CA, pp. 3-29). Transport can occur by a passive concentration-dependent mechanism, or can be linked to an energy source such as ATP hydrolysis or an ion gradient. Proteins that function in transport include carrier proteins, which bind to a specific solute and undergo a conformational change that translocates the bound solute across the membrane, and channel proteins, which form hydrophilic pores that allow specific solutes to diffuse through the membrane down an electrochemical solute gradient.

Carrier proteins which transport a single solute from one side of the membrane to the other are uniporters. In contrast, coupled transporters link the transfer of one solute with

25 simultaneous or sequential transfer of a second solute, either in the same direction (symport) or in the opposite direction (antiport). For example, intestinal and kidney epithelium contains a variety of symporter systems driven by the sodium gradient that exists across the plasma membrane. Sodium moves into the cell down its electrochemical gradient and brings the solute into the cell with it. The sodium gradient that provides the driving force for solute uptake is maintained by the ubiquitous  $\text{Na}^+/\text{K}^+$  ATPase system. Sodium-coupled transporters include the mammalian glucose transporter (SGLT1), iodide transporter (NIS), and multivitamin transporter (SMVT). All three transporters have twelve putative transmembrane segments, extracellular glycosylation sites, and cytoplasmically-oriented N- and C-termini. NIS plays a crucial role in the evaluation, diagnosis, and treatment of various thyroid pathologies because it is the molecular basis for radiiodide thyroid-imaging

techniques and for specific targeting of radioisotopes to the thyroid gland (Levy, O. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:5568-5573). SMVT is expressed in the intestinal mucosa, kidney, and placenta, and is implicated in the transport of the water-soluble vitamins, e.g., biotin and pantothenate (Prasad, P.D. et al. (1998) *J. Biol. Chem.* 273:7501-7506).

One of the largest families of transporters is the major facilitator superfamily (MFS), also called the uniporter-symporter-antiporter family. MFS transporters are single polypeptide carriers that transport small solutes in response to ion gradients. Members of the MFS are found in all classes of living organisms, and include transporters for sugars, oligosaccharides, phosphates, nitrates, nucleosides, monocarboxylates, and drugs. MFS transporters found in eukaryotes all have a structure comprising 12 transmembrane segments (Pao, S.S. et al. (1998) *Microbiol. Molec. Biol. Rev.* 62:1-34). The largest family of MFS transporters is the sugar transporter family, which includes the seven glucose transporters (GLUT1-GLUT7) found in humans that are required for the transport of glucose and other hexose sugars. These glucose transport proteins have unique tissue distributions and physiological functions. GLUT1 provides many cell types with their basal glucose requirements and transports glucose across epithelial and endothelial barrier tissues; GLUT2 facilitates glucose uptake or efflux from the liver; GLUT3 regulates glucose supply to neurons; GLUT4 is responsible for insulin-regulated glucose disposal; and GLUT5 regulates fructose uptake into skeletal muscle. Defects in glucose transporters are involved in a recently identified neurological syndrome causing infantile seizures and developmental delay, as well as glycogen storage disease, Fanconi-Bickel syndrome, and non-insulin-dependent diabetes mellitus (Mueckler, M. (1994) *Eur. J. Biochem.* 219:713-725; Longo, N. and L.J. Elsas (1998) *Adv. Pediatr.* 45:293-313).

Monocarboxylate anion transporters are proton-coupled symporters with a broad substrate specificity that includes L-lactate, pyruvate, and the ketone bodies acetate, acetoacetate, and beta-hydroxybutyrate. At least seven isoforms have been identified to date. The isoforms are predicted to have twelve transmembrane (TM) helical domains with a large intracellular loop between TM6 and TM7, and play a critical role in maintaining intracellular pH by removing the protons that are produced stoichiometrically with lactate during glycolysis. The best characterized H<sup>+</sup>-monocarboxylate transporter is that of the erythrocyte membrane, which transports L-lactate and a wide range of other aliphatic monocarboxylates. Other cells possess H<sup>+</sup>-linked monocarboxylate transporters with differing substrate and inhibitor selectivities. In particular, cardiac muscle and tumor cells have transporters that differ in their K<sub>m</sub> values for certain substrates, including stereoselectivity for L- over D-lactate, and in their sensitivity to inhibitors. There are Na<sup>+</sup>-monocarboxylate cotransporters on the luminal surface of intestinal and kidney epithelia, which allow the uptake of lactate, pyruvate, and ketone bodies in these tissues. In addition, there are specific and selective transporters for organic cations and organic anions

in organs including the kidney, intestine and liver. Organic anion transporters are selective for hydrophobic, charged molecules with electron-attracting side groups. Organic cation transporters, such as the ammonium transporter, mediate the secretion of a variety of drugs and endogenous metabolites, and contribute to the maintenance of intercellular pH (Poole, R.C. and A.P. Halestrap (1993) *Am. J. Physiol.* 264:C761-C782; Price, N.T. et al. (1998) *Biochem. J.* 329:321-328; and Martineau, K. and I. Haggstrom (1993) *J. Biotechnol.* 30:339-350).

ATP-binding cassette (ABC) transporters are members of a superfamily of membrane proteins that transport substances ranging from small molecules such as ions, sugars, amino acids, peptides, and phospholipids, to lipopeptides, large proteins, and complex hydrophobic drugs. ABC transporters consist of four modules: two nucleotide-binding domains (NBD), which hydrolyze ATP to supply the energy required for transport, and two membrane-spanning domains (MSD), each containing six putative transmembrane segments. These four modules may be encoded by a single gene, as is the case for the cystic fibrosis transmembrane regulator (CFTR), or by separate genes. When encoded by separate genes, each gene product contains a single NBD and MSD. These "half-molecules" form homo- and heterodimers, such as Tap1 and Tap2, the endoplasmic reticulum-based major histocompatibility (MHC) peptide transport system. Several genetic diseases are attributed to defects in ABC transporters, such as the following diseases and their corresponding proteins: cystic fibrosis (CFTR, an ion channel), adrenoleukodystrophy (adrenoleukodystrophy protein, ALDP), Zellweger syndrome (peroxisomal membrane protein-70, PMP70), and hyperinsulinemic hypoglycemia (sulfonylurea receptor, SUR). Overexpression of the multidrug resistance (MDR) protein, another ABC transporter, in human cancer cells makes the cells resistant to a variety of cytotoxic drugs used in chemotherapy (Taglicht, D. and S. Michaelis (1998) *Meth. Enzymol.* 292:130-162).

A number of metal ions such as iron, zinc, copper, cobalt, manganese, molybdenum, selenium, nickel, and chromium are important as cofactors for a number of enzymes. For example, copper is involved in hemoglobin synthesis, connective tissue metabolism, and bone development, by acting as a cofactor in oxidoreductases such as superoxide dismutase, ferroxidase (ceruloplasmin), and lysyl oxidase. Copper and other metal ions must be provided in the diet, and are absorbed by transporters in the gastrointestinal tract. Plasma proteins transport the metal ions to the liver and other target organs, where specific transporters move the ions into cells and cellular organelles as needed. Imbalances in metal ion metabolism have been associated with a number of disease states (Danks, D.M. (1986) *J. Med. Genet.* 23:99-106).

Transport of fatty acids across the plasma membrane can occur by diffusion, a high capacity, low affinity process. However, under normal physiological conditions a significant fraction of fatty acid transport appears to occur via a high affinity, low capacity protein-mediated transport process.

Fatty acid transport protein (FATP), an integral membrane protein with four transmembrane segments, is expressed in tissues exhibiting high levels of plasma membrane fatty acid flux, such as muscle, heart, and adipose. Expression of FATP is upregulated in 3T3-L1 cells during adipose conversion, and expression in COS7 fibroblasts elevates uptake of long-chain fatty acids (Hui, T. Y. et al. (1998) J. Biol. Chem. 273:27420-27429).

Mitochondrial carrier proteins are transmembrane-spanning proteins which transport ions and charged metabolites between the cytosol and the mitochondrial matrix. Examples include the ADP, ATP carrier protein; the 2-oxoglutarate/maleate carrier; the phosphate carrier protein; the pyruvate carrier; the dicarboxylate carrier which transports malate, succinate, fumarate, and phosphate; the tricarboxylate carrier which transports citrate and malate; and the Grave's disease carrier protein, a protein recognized by IgG in patients with active Grave's disease, an autoimmune disorder resulting in hyperthyroidism. Proteins in this family consist of three tandem repeats of an approximately 100 amino acid domain, each of which contains two transmembrane regions (Stryer, L. (1995) *Biochemistry*, W.H. Freeman and Company, New York NY, p. 551; PROSITE PDOC00189 Mitochondrial energy transfer proteins signature; Online Mendelian Inheritance in Man (OMIM) \*275000 Graves Disease).

This class of transporters also includes the mitochondrial uncoupling proteins, which create proton leaks across the inner mitochondrial membrane, thus uncoupling oxidative phosphorylation from ATP synthesis. The result is energy dissipation in the form of heat. Mitochondrial uncoupling proteins have been implicated as modulators of thermoregulation and metabolic rate, and have been proposed as potential targets for drugs against metabolic diseases such as obesity (Ricquier, D. et al. (1999) J. Int. Med. 245:637-642).

#### Ion Channels

The electrical potential of a cell is generated and maintained by controlling the movement of ions across the plasma membrane. The movement of ions requires ion channels, which form ion-selective pores within the membrane. There are two basic types of ion channels, ion transporters and gated ion channels. Ion transporters utilize the energy obtained from ATP hydrolysis to actively transport an ion against the ion's concentration gradient. Gated ion channels allow passive flow of an ion down the ion's electrochemical gradient under restricted conditions. Together, these types of ion channels generate, maintain, and utilize an electrochemical gradient that is used in 1) electrical impulse conduction down the axon of a nerve cell, 2) transport of molecules into cells against concentration gradients, 3) initiation of muscle contraction, and 4) endocrine cell secretion.

#### Ion Transporters

Ion transporters generate and maintain the resting electrical potential of a cell. Utilizing the energy derived from ATP hydrolysis, they transport ions against the ion's concentration gradient.

These transmembrane ATPases are divided into three families. The phosphorylated (P) class ion transporters, including Na<sup>+</sup>-K<sup>+</sup>-ATPase, Ca<sup>2+</sup>-ATPase, and H<sup>+</sup>-ATPase, are activated by a phosphorylation event. P-class ion transporters are responsible for maintaining resting potential distributions such that cytosolic concentrations of Na<sup>+</sup> and Ca<sup>2+</sup> are low and cytosolic concentration of K<sup>+</sup> is high. The vacuolar (V) class of ion transporters includes H<sup>+</sup> pumps on intracellular organelles, such as lysosomes and Golgi. V-class ion transporters are responsible for generating the low pH within the lumen of these organelles that is required for function. The coupling factor (F) class consists of H<sup>+</sup> pumps in the mitochondria. F-class ion transporters utilize a proton gradient to generate ATP from ADP and inorganic phosphate (P<sub>i</sub>).

The P-ATPases are hexamers of a 100 kD subunit with ten transmembrane domains and several large cytoplasmic regions that may play a role in ion binding (Scarborough, G. A. (1999) *Curr. Opin. Cell Biol.* 11:517-522). The V-ATPases are composed of two functional domains: the V<sub>1</sub> domain, a peripheral complex responsible for ATP hydrolysis, and the V<sub>0</sub> domain, an integral complex responsible for proton translocation across the membrane. The F-ATPases are structurally and evolutionarily related to the V-ATPases. The F-ATPase F<sub>0</sub> domain contains 12 copies of the c- subunit, a highly hydrophobic protein composed of two transmembrane domains and containing a single buried carboxyl group in TM2 that is essential for proton transport. The V-ATPase V<sub>0</sub> domain contains three types of homologous c subunits with four or five transmembrane domains and the essential carboxyl group in TM4 or TM3. Both types of complex also contain a subunit that may be involved in regulating the pH dependence of activity (Forger, M. (1999) J. Biol. Chem. 274:12951-12954).

The resting potential of the cell is utilized in many processes involving carrier proteins and gated ion channels. Carrier proteins utilize the resting potential to transport molecules into and out of the cell. Amino acid and glucose transport into many cells is linked to sodium ion co-transport (symport) so that the movement of Na<sup>+</sup> down an electrochemical gradient drives transport of the other molecule up a concentration gradient. Similarly, cardiac muscle links transfer of Ca<sup>2+</sup> out of the cell with transport of Na<sup>+</sup> into the cell (antiport).

#### Gated Ion Channels

Gated ion channels control ion flow by regulating the opening and closing of pores. The ability to control ion flux through various gating mechanisms allows ion channels to mediate such diverse signaling and homeostatic functions as neuronal and endocrine signaling, muscle contraction, fertilization, and regulation of ion and pH balance. Gated ion channels are categorized according to the manner of regulating the gating function. Mechanically-gated channels open their pores in response to mechanical stress; voltage-gated channels (e.g., Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, and Cl<sup>-</sup> channels) open their pores in response to changes in membrane potential; and ligand-gated channels (e.g.,

acetylcholine-, serotonin-, and glutamate-gated cation channels, and GABA- and glycine-gated chloride channels) open their pores in the presence of a specific ion, nucleotide, or neurotransmitter. The gating properties of a particular ion channel (i.e., its threshold for and duration of opening and closing) are sometimes modulated by association with auxiliary channel proteins and/or post translational modifications, such as phosphorylation.

5 Mechanically-gated or mechanosensitive ion channels act as transducers for the senses of touch, hearing, and balance, and also play important roles in cell volume regulation, smooth muscle contraction, and cardiac rhythm generation. A stretch-inactivated channel (SIC) was recently cloned from rat kidney. The SIC channel belongs to a group of channels which are activated by pressure or stress on the cell membrane and conduct both  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  (Suzuki, M. et al. (1999) *J. Biol. Chem.* 274:6330-6335).

The pore-forming subunits of the voltage-gated cation channels form a superfamily of ion channel proteins. The characteristic domain of these channel proteins comprises six transmembrane domains (S1-S6), a pore-forming region (P) located between S5 and S6, and intracellular amino and carboxy termini. In the  $\text{Na}^{+}$  and  $\text{Ca}^{2+}$  subfamilies, this domain is repeated four times, while in the  $\text{K}^{+}$  channel subfamily, each channel is formed from a tetramer of either identical or dissimilar subunits. The P region contains information specifying the ion selectivity for the channel. In the case of  $\text{K}^{+}$  channels, a GYG tripeptide is involved in this selectivity (Ishii, T.M. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:11651-11656).

20 Voltage-gated  $\text{Na}^{+}$  and  $\text{K}^{+}$  channels are necessary for the function of electrically excitable cells, such as nerve and muscle cells. Action potentials, which lead to neurotransmitter release and muscle contraction, arise from large, transient changes in the permeability of the membrane to  $\text{Na}^{+}$  and  $\text{K}^{+}$  ions. Depolarization of the membrane beyond the threshold level opens voltage-gated  $\text{Na}^{+}$  channels. Sodium ions flow into the cell, further depolarizing the membrane and opening more voltage-gated  $\text{Na}^{+}$  channels, which propagates the depolarization down the length of the cell. Depolarization also opens voltage-gated potassium channels. Consequently, potassium ions flow outward, which leads to repolarization of the membrane. Voltage-gated channels utilize charged residues in the fourth transmembrane segment (S4) to sense voltage change. The open state lasts only about 1 millisecond, at which time the channel spontaneously converts into an inactive state that cannot be opened irrespective of the membrane potential. Inactivation is mediated by the channel's N-terminus, which acts as a plug that closes the pore. The transition from an inactive to a closed state requires a return to resting potential.

Voltage-gated  $\text{Na}^{+}$  channels are heterotrimeric complexes composed of a 260 kDa pore-forming  $\alpha$  subunit that associates with two smaller auxiliary subunits,  $\beta 1$  and  $\beta 2$ . The  $\beta 2$  subunit is a integral

membrane glycoprotein that contains an extracellular Ig domain, and its association with  $\alpha$  and  $\beta 1$  subunits correlates with increased functional expression of the channel, a change in its gating properties, as well as an increase in whole cell capacitance due to an increase in membrane surface area (Isom, L.L. et al. (1995) *Cell* 83:433-442).

5 Non voltage-gated  $\text{Na}^{+}$  channels include the members of the amiloride-sensitive  $\text{Na}^{+}$  channel/degenerin (NaC/DEG) family. Channel subunits of this family are thought to consist of two transmembrane domains flanking a long extracellular loop, with the amino and carboxyl termini located within the cell. The NaC/DEG family includes the epithelial  $\text{Na}^{+}$  channel (ENaC) involved in  $\text{Na}^{+}$  reabsorption in epithelia including the airway, distal colon, cortical collecting duct of the kidney, and exocrine duct glands. Mutations in ENaC result in pseudohypoaldosteronism type 1 and Liddle's syndrome (pseudohypoaldosteronism). The NaC/DEG family also includes the recently characterized  $\text{H}^{+}$ -gated cation channels or acid-sensing ion channels (ASIC). ASIC subunits are expressed in the brain and form heteromultimeric  $\text{Na}^{+}$ -permeable channels. These channels require acid pH fluctuations for activation. ASIC subunits show homology to the degenerins, a family of mechanically-gated channels originally isolated from *C. elegans*. Mutations in the degenerins cause neurodegeneration. ASIC subunits may also have a role in neuronal function, or in pain perception, since tissue acidosis causes pain (Waldmann, R. and M. Lazdunski (1998) *Curr. Opin. Neurobiol.* 8:418-424; Eglen, R.M. et al. (1999) *Trends Pharmacol. Sci.* 20:337-342).

20  $\text{K}^{+}$  channels are located in all cell types, and may be regulated by voltage, ATP concentration, or second messengers such as  $\text{Ca}^{2+}$  and cAMP. In non-excitable tissue,  $\text{K}^{+}$  channels are involved in protein synthesis, control of endocrine secretions, and the maintenance of osmotic equilibrium across membranes. In neurons and other excitable cells, in addition to regulating action potentials and repolarizing membranes,  $\text{K}^{+}$  channels are responsible for setting resting membrane potential. The cytosol contains non-diffusible anions and, to balance this net negative charge, the cell contains a  $\text{Na}^{+}$ .  $\text{K}^{+}$  pump and ion channels that provide the redistribution of  $\text{Na}^{+}$ ,  $\text{K}^{+}$ , and  $\text{Cl}^{-}$ . The pump actively transports  $\text{Na}^{+}$  out of the cell and  $\text{K}^{+}$  into the cell in a 3:2 ratio. Ion channels in the plasma membrane allow  $\text{K}^{+}$  and  $\text{Cl}^{-}$  to flow by passive diffusion. Because of the high negative charge within the cytosol,  $\text{Cl}^{-}$  flows out of the cell. The flow of  $\text{K}^{+}$  is balanced by an electromotive force pulling  $\text{K}^{+}$  into the cell, and a  $\text{K}^{+}$  concentration gradient pushing  $\text{K}^{+}$  out of the cell. Thus, the resting membrane potential is primarily regulated by  $\text{K}^{+}$  flow (Salhoffs, L. and T. Jegla (1995) *Neuron* 15:489-492).

30 Potassium channel subunits of the Shaker-like superfamily all have the characteristic six transmembrane/1 pore domain structure. Four subunits combine as homo- or heterotetramers to form functional K channels. These pore-forming subunits also associate with various cytoplasmic  $\beta$  subunits that alter channel inactivation kinetics. The Shaker-like channel family includes the voltage-

gated K<sup>+</sup> channels as well as the delayed rectifier type channels such as the human ether-a-go-go related gene (HERG) associated with long QT, a cardiac dysrhythmia syndrome (Curran, M.E. (1998) *Curr. Opin. Biotechnol.* 9:565-572; Kaczorowski, G.J. and M.L. Garcia (1999) *Curr. Opin. Chem. Biol.* 3:448-458).

5 A second superfamily of K<sup>+</sup> channels is composed of the inward rectifying channels (Kir).

Kir channels have the property of preferentially conducting K<sup>+</sup> currents in the inward direction. These proteins consist of a single potassium selective pore domain and two transmembrane domains, which correspond to the fifth and sixth transmembrane domains of voltage-gated K<sup>+</sup> channels. Kir subunits also associate as tetramers. The Kir family includes ROMK1, mutations in which lead to Bartter syndrome, a renal tubular disorder. Kir channels are also involved in regulation of cardiac pacemaker activity, seizures and epilepsy, and insulin regulation (Doupnik, C.A. et al. (1995) *Curr. Opin. Neurobiol.* 5:268-277; Curran, *EMBO*).

The recently recognized TWIK K<sup>+</sup> channel family includes the mammalian TWIK-1, TREK-1 and TASK proteins. Members of this family possess an overall structure with four transmembrane domains and two P domains. These proteins are probably involved in controlling the resting potential in a large set of cell types (Duprat, F. et al. (1997) *EMBO J* 16:5464-5471).

The voltage-gated Ca<sup>2+</sup> channels have been classified into several subtypes based upon their electrophysiological and pharmacological characteristics. L-type Ca<sup>2+</sup> channels are predominantly expressed in heart and skeletal muscle where they play an essential role in excitation-contraction coupling. T-type channels are important for cardiac pacemaker activity, while N-type and P/Q-type channels are involved in the control of neurotransmitter release in the central and peripheral nervous system. The L-type and N-type voltage-gated Ca<sup>2+</sup> channels have been purified and, though their functions differ dramatically, they have similar subunit compositions. The channels are composed of three subunits. The  $\alpha_1$  subunit forms the membrane pore and voltage sensor, while the  $\alpha_2\delta$  and  $\beta$  subunits modulate the voltage-dependence, gating properties, and the current amplitude of the channel. These subunits are encoded by at least six  $\alpha_1$ , one  $\alpha_2\delta$ , and four  $\beta$  genes. A fourth subunit,  $\gamma$ , has been identified in skeletal muscle (Walker, D. et al. (1998) *J. Biol. Chem.* 273:2361-2367; McCleskey, E.W. (1994) *Curr. Opin. Neurobiol.* 4:304-312).

The transient receptor family (Trp) of calcium ion channels are thought to mediate capacitative calcium entry (CCE). CCE is the Ca<sup>2+</sup> influx into cells to resupply Ca<sup>2+</sup> stores depleted by the action of inositol triphosphate (IP3) and other agents in response to numerous hormones and growth factors. Trp and Trp-like were first cloned from *Drosophila* and have similarity to voltage gated Ca<sup>2+</sup> channels in the S3 through S6 regions. This suggests that Trp and/or related proteins may form mammalian CCC entry channels (Zhu, X. et al. (1996) *Cell* 85:661-671; Boulay, G. et al. (1997) *J. Biol. Chem.*

27229672-29680). Melastatin is a gene isolated in both the mouse and human, and whose expression in melanoma cells is inversely correlated with melanoma aggressiveness *in vivo*. The human CDNA transcript corresponds to a 1533-amino acid protein having homology to members of the Trp family. It has been proposed that the combined use of melastatin mRNA expression status and tumor thickness might allow for the determination of subgroups of patients at both low and high risk for developing metastatic disease (Duncan, L.M. et al (2001) *J. Clin. Oncol.* 19:568-576).

Chloride channels are necessary in endocrine secretion and in regulation of cytosolic and organelle pH. In secretory epithelial cells, Cl<sup>-</sup> enters the cell across a basolateral membrane through an Na<sup>+</sup>, K<sup>+</sup>/Cl<sup>-</sup> cotransporter, accumulating in the cell above its electrochemical equilibrium concentration.

10 Secretion of Cl<sup>-</sup> from the apical surface, in response to hormonal stimulation, leads to flow of Na<sup>+</sup> and water into the secretory lumen. The cystic fibrosis transmembrane conductance regulator (CFTR) is a chloride channel encoded by the gene for cystic fibrosis, a common fatal genetic disorder in humans. CFTR is a member of the ABC transporter family, and is composed of two domains each consisting of six transmembrane domains followed by a nucleotide-binding site. Loss of CFTR function decreases transepithelial water secretion and, as a result, the layers of mucus that coat the respiratory tree, pancreatic ducts, and intestine are dehydrated and difficult to clear. The resulting blockage of these sites leads to pancreatic insufficiency, "meconium ileus", and devastating "chronic obstructive pulmonary disease" (Al-Awqali, Q. et al. (1992) *J. Exp. Biol.* 172:245-260).

The voltage-gated chloride channels (ClC) are characterized by 10-12 transmembrane domains, as well as two small globular domains known as CBS domains. The ClC subunits probably function as homotetramers. ClC proteins are involved in regulation of cell volume, membrane potential stabilization, signal transduction, and transepithelial transport. Mutations in ClC-1, expressed predominantly in skeletal muscle, are responsible for autosomal recessive generalized myotonia and autosomal dominant myotonia congenita, while mutations in the kidney channel ClC-5 lead to kidney stones (Jeansch, T.J. (1996) *Curr. Opin. Neurobiol.* 6:303-310).

Ligand-gated channels open their pores when an extracellular or intracellular mediator binds to the channel. Neurotransmitter-gated channels are channels that open when a neurotransmitter binds to their extracellular domain. These channels exist in the postsynaptic membrane of nerve or muscle cells. There are two types of neurotransmitter-gated channels. Sodium channels open in response to excitatory neurotransmitters, such as acetylcholine, glutamate, and serotonin. This opening causes an influx of Na<sup>+</sup> and produces the initial localized depolarization that activates the voltage-gated channels and starts the action potential. Chloride channels open in response to inhibitory neurotransmitters, such as  $\gamma$ -aminobutyric acid (GABA) and glycine, leading to hyperpolarization of the membrane and the subsequent generation of an action potential. Neurotransmitter-gated ion channels have four

transmembrane domains and probably function as pentamers (Jentsch, *supra*). Amino acids in the second transmembrane domain appear to be important in determining channel permeation and selectivity (Sather, W.A. et al. (1994) *Curr. Opin. Neurobiol.* 4:313-323).

Ligand-gated channels can be regulated by intracellular second messengers. For example, calcium-activated K<sup>+</sup> channels are gated by internal calcium ions. In nerve cells, an influx of calcium during depolarization opens K<sup>+</sup> channels to modulate the magnitude of the action potential (Ishi et al., *supra*). The large conductance (BK) channel has been purified from brain and its subunit composition determined. The  $\alpha$  subunit of the BK channel has seven rather than six transmembrane domains in contrast to voltage-gated K<sup>+</sup> channels. The extra transmembrane domain is located at the subunit N-terminus. A 28-amino-acid stretch in the C-terminal region of the subunit (the "calcium bowl" region) contains many negatively charged residues and is thought to be the region responsible for calcium binding. The  $\beta$  subunit consists of two transmembrane domains connected by a glycosylated extracellular loop, with intracellular N- and C-termini (Kaczorowski, *supra*; Vergara, C. et al. (1998) *Curr. Opin. Neurobiol.* 8:321-329).

Cyclic nucleotide-gated (CNG) channels are gated by cytosolic cyclic nucleotides. The best examples of these are the cAMP-gated Na<sup>+</sup> channels involved in olfaction and the cGMP-gated cation channels involved in vision. Both systems involve ligand-mediated activation of a G-protein coupled receptor which then alters the level of cyclic nucleotide within the cell. CNG channels also represent a major pathway for Ca<sup>2+</sup> entry into neurons, and play roles in neuronal development and plasticity. CNG channels are tetramers containing at least two types of subunits, an  $\alpha$  subunit which can form functional homomeric channels, and a  $\beta$  subunit, which modulates the channel properties. All CNG subunits have six transmembrane domains and a pore forming region between the fifth and sixth transmembrane domains, similar to voltage-gated K<sup>+</sup> channels. A large C-terminal domain contains a cyclic nucleotide binding domain, while the N-terminal domain confers variation among channel subtypes (Zuñali, F. et al. (1997) *Curr. Opin. Neurobiol.* 7:404-412).

The activity of other types of ion channel proteins may also be modulated by a variety of intracellular signalling proteins. Many channels have sites for phosphorylation by one or more protein kinases including protein kinase A, protein kinase C, tyrosine kinase, and casein kinase II, all of which regulate ion channel activity in cells. Kir channels are activated by the binding of the G $\beta\gamma$  subunits of heterotrimeric G-proteins (Reimann, F. and F.M. Ashcroft (1999) *Curr. Opin. Cell. Biol.* 11:503-508). Other proteins are involved in the localization of ion channels to specific sites in the cell membrane. Such proteins include the PDZ domain proteins known as MAGUKs (membrane-associated guanylate kinases) which regulate the clustering of ion channels at neuronal synapses (Craven, S.E. and D.S. Bredt (1998) *Cell* 93:495-498).

# Disease Correlation

The etiology of numerous human diseases and disorders can be attributed to defects in the transport of molecules across membranes. Defects in the trafficking of membrane-bound transporters and ion channels are associated with several disorders, e.g., cystic fibrosis, glucose-galactose

malabsorption syndrome, hypercholesterolemia, von Gierke disease, and certain forms of diabetes mellitus. Single-gene defect diseases resulting in an inability to transport small molecules across membranes include, e.g., cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease (van't Hoff, W.G. (1996) *Exp. Nephrol.* 4:253-262; Talente, G.M. et al. (1994) *Ann. Intern. Med.* 120:218-226; and Chillon, M. et al. (1995) *New Engl. J. Med.* 332:1475-1480).

Human diseases caused by mutations in ion channel genes include disorders of skeletal muscle, cardiac muscle, and the central nervous system. Mutations in the pore-forming subunits of sodium and chloride channels cause myotonia, a muscle disorder in which relaxation after voluntary contraction is delayed. Sodium channel myotonias have been treated with channel blockers. Mutations in muscle sodium and calcium channels cause forms of periodic paralysis, while mutations in the sarcoplasmic calcium release channel, T-tubule calcium channel, and muscle sodium channel cause malignant hyperthermia. Cardiac arrhythmia disorders such as the long QT syndromes and idiopathic ventricular fibrillation are caused by mutations in potassium and sodium channels (Cooper, E.C. and L.Y. Jan (1998) *Proc. Natl. Acad. Sci. USA* 96:4759-4766). All four known human idiopathic epilepsy genes code for ion channel proteins (Berkovic, S.F. and I.E. Scheffer (1999) *Curr. Opin. Neurology* 12:177-182). Other neurological disorders such as ataxias, hemiplegic migraine and hereditary deafness can also result from mutations in ion channel genes (Jen, J. (1999) *Curr. Opin. Neurobiol.* 9:274-280; Cooper, *supra*).

Ion channels have been the target for many drug therapies. Neurotransmitter-gated channels have been targeted in therapies for treatment of insomnia, anxiety, depression, and schizophrenia. Voltage-gated channels have been targeted in therapies for arrhythmia, ischemic stroke, head trauma, and neurodegenerative disease (Taylor, C.P. and L.S. Narasimhan (1997) *Adv. Pharmacol.* 39:47-98). Various classes of ion channels also play an important role in the perception of pain, and thus are potential targets for new analgesics. These include the vanilloid-gated ion channels, which are activated by the vanilloid capsaicin, as well as by noxious heat. Local anesthetics such as lidocaine and mexiletine which block voltage-gated Na<sup>+</sup> channels have been useful in the treatment of neuropathic pain (Eglen, *supra*).

Ion channels in the immune system have recently been suggested as targets for immunomodulation. T-cell activation depends upon calcium signaling, and a diverse set of T-cell specific ion channels has been characterized that affect this signaling process. Channel blocking agents

can inhibit secretion of lymphokines, cell proliferation, and killing of target cells. A peptide antagonist of the T-cell potassium channel Kv1.3 was found to suppress delayed-type hypersensitivity and allogeneic responses in pigs, validating the idea of channel blockers as safe and efficacious immunosuppressants (Cohnan, M.D. and K.G. Chandry (1997) *Curr. Opin. Biotechnol.* 8:749-756).

The discovery of new transporters and ion channels, and the polynucleotides encoding them, satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of transport, neurological, muscle, immunological, and cell proliferative disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of transporters and ion channels.

#### SUMMARY OF THE INVENTION

The invention features purified polypeptides, transporters and ion channels, referred to collectively as "TRICH" and individually as "TRICH-1," "TRICH-2," "TRICH-3," "TRICH-4," "TRICH-5," "TRICH-6," "TRICH-7," "TRICH-8," "TRICH-9," "TRICH-10," "TRICH-11," "TRICH-12," "TRICH-13," "TRICH-14," "TRICH-15," "TRICH-16," "TRICH-17," "TRICH-18," "TRICH-19," "TRICH-20," "TRICH-21," "TRICH-22," "TRICH-23," "TRICH-24," "TRICH-25," "TRICH-26," "TRICH-27," "TRICH-28," "TRICH-29," "TRICH-30," "TRICH-31," and "TRICH-32." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-32.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-32. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:33-64.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional TRICH, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and e) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and f) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) combining the polypeptide with at least one test compound under

suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

- The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-32. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

- The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:33-64, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

- The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, iii) a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, iv) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:33-64, iii) a polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the

- polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

#### BRIEF DESCRIPTION OF THE TABLES

- Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the present invention.

- Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

- Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

- Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

- Table 5 shows the representative cDNA library for polynucleotides of the invention.

- Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

- Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

#### DESCRIPTION OF THE INVENTION

- Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

- It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a

reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

#### DEFINITIONS

"TRICH" refers to the amino acid sequences of substantially purified TRICH obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of TRICH. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TRICH either by directly interacting with TRICH or by acting on components of the biological pathway in which TRICH participates.

An "allelic variant" is an alternative form of the gene encoding TRICH. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding TRICH include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as TRICH or a polypeptide with at least one functional characteristic of TRICH. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding TRICH, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding TRICH. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent TRICH. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge,

solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of TRICH is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of TRICH. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TRICH either by directly interacting with TRICH or by acting on components of the biological pathway in which TRICH participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind TRICH polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA, RNA, peptide nucleic acid (PNA), oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethylphosphonates; or oligonucleotides having modified bases such as 5-methyl cytosine, 2-deoxyuracil, or 7-deaza-2-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic TRICH, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-ACT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding TRICH or fragments of TRICH may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate, SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCC, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

Original Residue	Conservative Substitution
Ala	Gly, Ser
Arg	His, Lys
Asn	Asp, Gln, His
Asp	Asn, Glu
Cys	Ala, Ser
Gln	Asn, Glu, His
Glu	Asp, Gln, His
Gly	Ala
His	Asn, Arg, Gln, Glu
Ile	Leu, Val
Leu	Ile, Val
Lys	Arg, Gln, Glu
Met	Leu, Ile
Phe	His, Met, Leu, Trp, Tyr
Ser	Cys, Thr
Thr	Ser, Val
Trp	Phe, Tyr
Tyr	His, Phe, Trp
Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated, or decreased, downregulated, or

absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

A "fragment" is a unique portion of TRICH or the polynucleotide encoding TRICH which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:33-64 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:33-64, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:33-64 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:33-64 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:33-64 and the region of SEQ ID NO:33-64 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-32 is encoded by a fragment of SEQ ID NO:33-64. A fragment of SEQ ID NO:1-32 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-32. For example, a fragment of SEQ ID NO:1-32 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-32. The precise length of a fragment of SEQ ID NO:1-32 and the region of SEQ ID NO:1-32 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to

the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore, achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191.

For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The

"BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12a sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April 21-2000) with biasp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for

24

example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity.

Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY, specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour.

25

Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, steared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C<sub>4</sub> or R<sub>4</sub> analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of TRICH which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of TRICH which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of TRICH. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of TRICH.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or

synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an TRICH may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of TRICH.

"Probe" refers to nucleic acid sequences encoding TRICH, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR

Primer3: A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

- 5 Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PRIMO primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimerGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences.
- 20 Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

- 25 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *et al.* The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

- 5 A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

- 10 "Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

- 15 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing TRICH, nucleic acids encoding TRICH, or fragments thereof may comprise a bodily fluid, an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate, a tissue; a tissue print; etc.

- 20 The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.
- 25 The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

- 30 A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.
- "Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers,

microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

5 "Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

10 A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), SUTDA.

25 A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant

identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule.

Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

10 A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

## THE INVENTION

20 The invention is based on the discovery of new human transporters and ion channels (TRICH), the polynucleotides encoding TRICH, and the use of these compositions for the diagnosis, treatment, or prevention of transport, neurological, muscle, immunological, and cell proliferative disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO.) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO.) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

30 Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO.) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO.) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank

homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO.) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are transporters and ion channels. For example, SEQ ID NO.5 is 83% identical to rat GABA receptor rho-3 subunit precursor (GenBank ID g1060975) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.7e-206, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO.5 also contains a neurotransmitter-gated ion channel domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO.5 is a neurotransmitter-gated ion channel. In an alternate example, SEQ ID NO.16 is 57% identical to human Na<sup>+</sup>/glucose cotransporter (GenBank ID g338055) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 2.4e-181, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO.16 also contains a sodium:olute symporter family domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO.16 is a Na<sup>+</sup>/glucose cotransporter. In an alternate example, SEQ ID NO.27 is 53% identical to human ATP-binding cassette transporter-1 (ABC-1) (GenBank ID g4128033) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 0.0, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO.27 also contains an ABC transporter domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from

BLIMPS, MOTIFS, and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO.27 is an ABC transporter. In an alternate example, SEQ ID NO.12 is 45% identical to rat thyroid sodium/iodide symporter NIS (GenBank ID g1399954) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 3.0e-143, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO.12 also contains a sodium:olute symporter family domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO.12 is a sodium:olute symporter. SEQ ID NO.14, SEQ ID NO.6-11, SEQ ID NO.13-15, SEQ ID NO.17-26, and SEQ ID NO.28-32 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO.1-32 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO.) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO.33-64 or that distinguish between SEQ ID NO.33-64 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblies comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 6724643H1 is the identification number of an Incyte cDNA sequence, and LUN1.TM101 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71495515V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g5746200) which contributed to the assembly of the full length polynucleotide sequences. In addition, the identification numbers in column 5 may identify sequences derived from the ENSEMBL (The Sanger Centre, Cambridge, UK) database (i.e., those sequences

including the designation "ENST"). Alternatively, the identification numbers in column 5 may be derived from the NCBI RefSeq Nucleotide Sequence Records Database (*i.e.*, those sequences including the designation "NM" or "NT") or the NCBI RefSeq Protein Sequence Records (*i.e.*, those sequences including the designation "NP"). Alternatively, the identification numbers in column 5 may refer to assemblies of both cDNA and GenScan-predicted exons brought together by an "exon stitching" algorithm. For example, FL\_XXXXXX\_N<sub>1</sub>\_YYYYY\_N<sub>2</sub> represents a "stitched" sequence in which XXXXXX is the identification number of the cluster of sequences to which the algorithm was applied, and YYYYY is the number of the prediction generated by the algorithm, and N<sub>1</sub>, N<sub>2</sub>, if present, represent specific exons that may have been manually edited during analysis (See Example V).

Alternatively, the identification numbers in column 5 may refer to assemblies of exons brought together by an "exon-stretching" algorithm. For example, FL\_XXXXXX\_gAAAAA\_gBBBBB\_1\_N is the identification number of a "stretched" sequence, with XXXXXX being the Incyte project identification number, gAAAAA being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, gBBBBB being the GenBank identification number or NCBI RefSeq identification number of the nearest GenBank protein homolog, and N referring to specific exons (See Example V). In instances where a RefSeq sequence was used as a protein homolog for the "exon-stretching" algorithm, a RefSeq identifier (denoted by "NM," "NP," or "NT") may be used in place of the GenBank identifier (*i.e.*, gBBBBB).

Alternatively, a prefix identifies component sequences that were hand-edited, predicted from genomic DNA sequences, or derived from a combination of sequence analysis methods. The following Table lists examples of component sequence prefixes and corresponding sequence analysis methods associated with the prefixes (see Example IV and Example V).

Prefix	Type of analysis and/or examples of programs
GNN, GFG, ENST	Exon prediction from genomic sequences using, for example, GENSCAN (Stanford University, CA, USA) or FGES (Computer Genomics Group, The Sanger Centre, Cambridge, UK).
GBI	Hand-edited analysis of genomic sequences.
FL	Stitched or stretched genomic sequences (see Example V).

In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte

cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses TRICH variants. A preferred TRICH variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the TRICH amino acid sequence, and which contains at least one functional or structural characteristic of TRICH.

The invention also encompasses polynucleotides which encode TRICH. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:33-64, which encodes TRICH. The polynucleotide sequences of SEQ ID NO:33-64, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding TRICH. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding TRICH. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:33-64 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:33-64. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of TRICH.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding TRICH, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring TRICH, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode TRICH and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring TRICH under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding TRICH or its derivatives possessing a substantially different codon usage, *e.g.*, inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide

occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding TRICH and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode TRICH and TRICH derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding TRICH or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:33-64 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) *Short Protocols in Molecular Biology*, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) *Molecular Biology and Biotechnology*, Wiley VCH, New York NY, pp. 836-833.)

The nucleic acid sequences encoding TRICH may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic

DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.)

Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Trigila, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode TRICH may be cloned in recombinant DNA molecules that direct expression of TRICH, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of

the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express TRICH.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter TRICH-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULAR BREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) *Nat. Biotechnol.* 17:793-797; Christians, F.C. et al. (1999) *Nat. Biotechnol.* 17:259-264; and Cramer, A. et al. (1996) *Nat. Biotechnol.* 14:315-319) to alter or improve the biological properties of TRICH, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding TRICH may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Curuthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, TRICH itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) *Proteins, Structures and Molecular Properties*, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of TRICH, or any part thereof, may be altered during direct synthesis and/or combined with sequences

from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chitez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:397-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, *supra*, pp. 28-53.)

In order to express a biologically active TRICH, the nucleotide sequences encoding TRICH or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding TRICH. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding TRICH. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding TRICH and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Schurf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding TRICH and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. (See, e.g., Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding TRICH. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or

- animal cell systems. (See, e.g., Sambrook, *SUBP*; Ausubel, *SUBP*; Van Hecke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Butler, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Sonja (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding TRICH. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding TRICH can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORI1 plasmid (Life Technologies). Ligation of sequences encoding TRICH into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Hecke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of TRICH are needed, e.g., for the production of antibodies, vectors which direct high level expression of TRICH may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

- Yeast expression systems may be used for production of TRICH. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *SUBP*; Butler, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Sorey, C.A. et al. (1994) *BioTechnology* 12:181-184.)

Plant systems may also be used for expression of TRICH. Transcription of sequences encoding TRICH may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311).

- Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding TRICH may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses TRICH in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

- Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of TRICH in cell lines is preferred. For example, sequences encoding TRICH can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

- Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *aprt* cells, respectively. (See, e.g., Wigler, M. et al. (1977) *Cell* 11:223-232; Lowy, I. et al. (1980) *Cell* 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *uid* and *pat* confer resistance to chloroform and phosphotriazin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980)

Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding TRICH is inserted within a marker gene sequence, transformed cells containing sequences encoding TRICH can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding TRICH under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding TRICH and that express TRICH may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of TRICH using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on TRICH is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Colligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunotechnical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding TRICH include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding TRICH, or any fragments thereof, may be cloned into a vector for

the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding TRICH may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode TRICH may be designed to contain signal sequences which direct secretion of TRICH through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding TRICH may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric TRICH protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of TRICH activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, c-myc, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, c-myc, and hemagglutinin (HA) enable immunofluorescence purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize

these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the TRICH encoding sequence and the heterologous protein sequence, so that TRICH may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *SUPRA*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled TRICH may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T<sub>7</sub>, T<sub>3</sub>, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

TRICH of the present invention or fragments thereof may be used to screen for compounds that specifically bind to TRICH. At least one and up to a plurality of test compounds may be screened for specific binding to TRICH. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of TRICH, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) *Current Protocols in Immunology* 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which TRICH binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express TRICH, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing TRICH or cell membrane fractions which contain TRICH are then contacted with a test compound and binding, stimulation, or inhibition of activity of either TRICH or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with TRICH, either in solution or affixed to a solid support, and detecting the binding of TRICH to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

TRICH of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of TRICH. Such compounds may include agonists, antagonists, or partial

or inverse agonists. In one embodiment, an assay is performed under conditions permissive for TRICH activity, wherein TRICH is combined with at least one test compound, and the activity of TRICH in the presence of a test compound is compared with the activity of TRICH in the absence of the test compound. A change in the activity of TRICH in the presence of the test compound is indicative of a compound that modulates the activity of TRICH. Alternatively, a test compound is combined with an

in *vitro* or cell-free system comprising TRICH under conditions suitable for TRICH activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of TRICH may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding TRICH or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,357.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest

disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (*neo*; Capechi, M.R. (1989) *Science* 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the *Cre-loxP* system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marrn, J.D. (1996) *Clin. Invest.* 97:1999-2002; Wagner, K.U. et al. (1997) *Nucleic Acids Res.* 25:4323-4330).

Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding TRICH may also be manipulated *in vitro* in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) *Science* 282:1145-1147).

Polynucleotides encoding TRICH can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding TRICH is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with

potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress TRICH, e.g., by secreting TRICH in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

# THERAPEUTICS

5 Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of TRICH and transporters and ion channels. In addition, the expression of TRICH is closely associated with adrenal, testicular, and prostate tumors, Crohn's disease, teratocarcinoma and dendritic cells, brain, lung, ileum, small intestine, uterine myometrial, colon, and pancreatic tissues. Therefore, TRICH appears to play a role in transport, neurological, muscle, immunological, and cell proliferative disorders. In the treatment of disorders associated with increased TRICH expression or activity, it is desirable to decrease the expression or activity of TRICH. In the treatment of disorders associated with decreased TRICH expression or activity, it is desirable to increase the expression or activity of TRICH.

10 Therefore, in one embodiment, TRICH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, cataplexy, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis, neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia, and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other

extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebellofudinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, cataplexy, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a muscle disorder such as cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, pheochromocytoma, and myopathies including encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, ophthalmoplegia, and acid maltase deficiency (AMD, also known as Pompe's disease); an immunological disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune thyroiditis, polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxicity, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hyper eosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal

circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma, and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

10 In another embodiment, a vector capable of expressing TRICH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those described above.

15 In a further embodiment, a composition comprising a substantially purified TRICH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of TRICH may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TRICH including, but not limited to, those listed above.

20 In a further embodiment, an antagonist of TRICH may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of TRICH. Examples of such disorders include, but are not limited to, those transport, neurological, muscle, immunological, and cell proliferative disorders described above. In one aspect, an antibody which specifically binds TRICH may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express TRICH.

25 In an additional embodiment, a vector expressing the complement of the polynucleotide encoding TRICH may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of TRICH including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of TRICH may be produced using methods which are generally known in the art. In particular, purified TRICH may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind TRICH. Antibodies to TRICH may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with TRICH or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolcedin, pluronic polyols, polyamides, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guérin) and *Corynebacterium parvum* are especially preferable.

15 It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to TRICH have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of TRICH amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

20 Monoclonal antibodies to TRICH may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cole, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and

25 Cole S.P. et al. (1984) *Mol. Cell Biol.* 4:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce TRICH-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be

generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for TRICH may also be generated. For example, such fragments include, but are not limited to, F(ab)<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab)<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between TRICH and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering TRICH epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for TRICH. Affinity is expressed as an association constant, K<sub>a</sub>, which is defined as the molar concentration of TRICH-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K<sub>a</sub> determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple TRICH epitopes, represents the average affinity, or avidity, of the antibodies for TRICH. The K<sub>a</sub> determined for a preparation of monoclonal antibodies, which are monospecific for a particular TRICH epitope, represents a true measure of affinity. High-affinity antibody preparations with K<sub>a</sub> ranging from about 10<sup>8</sup> to 10<sup>12</sup> L/mole are preferred for use in immunoassays in which the TRICH-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K<sub>a</sub> ranging from about 10<sup>6</sup> to 10<sup>7</sup> L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of TRICH, preferably in active form, from the antibody (Catty, D. (1988) *Antibodies, Volume I: A Practical Approach*, IRL Press, Washington DC; Liddle, J.E. and A. Cryer (1991) *A Practical Guide to Monoclonal Antibodies*, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of TRICH-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding TRICH, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding TRICH. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding TRICH. (See, e.g., Agrawal, S., ed. (1996) *Antisense Therapeutics*, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Sculnon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uekert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding TRICH may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaise, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial

hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Sonnia (1997) *Nature* 389:239-242)). (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as *Candida albicans* and *Paracoccidioides brasiliensis*, and protozoan parasites such as *Plasmodium falciparum* and *Toxoplasma cruzi*). In the case where a genetic deficiency in TRICH expression or regulation causes disease, the expression of TRICH from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in TRICH are treated by constructing mammalian expression vectors encoding TRICH and introducing these vectors by mechanical means into TRICH-deficient cells. Mechanical transfer technologies for use with cells *in vivo* or *ex vivo* include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J.-L. and H. Récapon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of TRICH include, but are not limited to, the pCDNA 3.1, EPTTAG, PROCMV, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OPF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). TRICH may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Biau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-Rex plasmid (Invitrogen)); the octadecane-inducible promoter (available in the plasmids PVGGRXR and PIND; Invitrogen); the FK506/fapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Biau, H.M. *supra*), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding TRICH from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver

polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to TRICH expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding TRICH under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PTB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an

appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zulfikery, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Riggs ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference.

Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bomyhad, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding TRICH to cells which have one or more genetic abnormalities with respect to the expression of TRICH. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact sites in the pancreas (Csele, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinuzzi, P.A. et al. (1999) *Annu.*

Rev. Nutr. 19:511-544 and Verma, I.M. and N. Sonnia (1997) Nature 18:389-239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding TRICH to target cells which have one or more genetic abnormalities with respect to the expression of TRICH. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing TRICH to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goinis, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding TRICH to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for TRICH into the alphavirus genome in place of the capsid-coding region results in the production of a large number of TRICH-coding RNAs and the synthesis of high levels of TRICH in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application

(Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of TRICH into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding TRICH.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding TRICH. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2'-O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of RNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding TRICH.

- 10 Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased TRICH expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding TRICH may be therapeutically useful, and in the treatment of disorders associated with decreased TRICH expression or activity, a compound which specifically promotes expression of the polynucleotide encoding TRICH may be therapeutically useful.

- 20 At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding TRICH is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding TRICH are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding TRICH. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of

the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Brutce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Brutce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

- 10 Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vitro, in vivo, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

- 20 An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Mack Publishing, Easton PA). Such compositions may consist of TRICH, antibodies to TRICH, and mimetics, agonists, antagonists, or inhibitors of TRICH.

- 25 The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

- 30 Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S.

Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising TRICH or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, TRICH or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) *Science* 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example TRICH or fragments thereof, antibodies of TRICH, and agonists, antagonists or inhibitors of TRICH, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population) or LD<sub>50</sub> (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD<sub>50</sub>/ED<sub>50</sub> ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED<sub>50</sub> with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy.

Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 µg to 100,000 µg, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

# DIAGNOSTICS

In another embodiment, antibodies which specifically bind TRICH may be used for the diagnosis of disorders characterized by expression of TRICH, or in assays to monitor patients being treated with TRICH or agonists, antagonists, or inhibitors of TRICH. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for TRICH include methods which utilize the antibody and a label to detect TRICH in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring TRICH, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of TRICH expression. Normal or standard values for TRICH expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to TRICH under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of TRICH expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding TRICH may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of TRICH may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of TRICH, and to monitor regulation of TRICH levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding TRICH or closely related molecules may be used to identify nucleic acid sequences which encode TRICH. The specificity of the probe, whether it is made

from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding TRICH, allelic variants, or related sequences.

5 Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the TRICH encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:33-64 or from genomic sequences including promoters, enhancers, and introns of the TRICH gene.

10 Means for producing specific hybridization probes for DNAs encoding TRICH include the cloning of polynucleotide sequences encoding TRICH or TRICH derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA

15 polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as <sup>32</sup>P or <sup>35</sup>S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding TRICH may be used for the diagnosis of disorders

associated with expression of TRICH. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, cataplexy, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g.,

angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis,

25 cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis, neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychosis, and schizophrenia, and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease,

30 cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Meckles disease, occipital horn syndrome, von Gierke disease, cystinuria, immunoglobulinuria, Hartup disease, and

Fanconi disease, a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalofibrogenal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD),

15 akathisia, amnesia, cataplexy, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia, a muscle disorder such as cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, Becker's muscular dystrophy, myotonic dystrophy, central core disease, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, infectious myositis, polymyositis, dermatomyositis, inclusion body myositis, thyrotoxic myopathy, ethanol myopathy, angina, anaphylactic shock, arrhythmias, asthma, cardiovascular shock, Cushing's syndrome, hypertension, hypoglycemia, myocardial infarction, migraine, pheochromocytoma, and myopathies including encephalopathy, epilepsy, Kearns-Sayre syndrome, lactic acidosis, myoclonic disorder, ophthalmoplegia, and acid malase deficiency (AMD, also known as Pompe's disease); an immunological disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune

25 polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxicins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypercosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic

30

anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding TRICH may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered TRICH expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding TRICH may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding TRICH may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding TRICH in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of TRICH, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding TRICH, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding TRICH may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding TRICH, or a fragment of a polynucleotide complementary to the polynucleotide encoding TRICH, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding TRICH may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding TRICH are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplicons in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the

alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of TRICH include radiolabeling or blotting nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, TRICH fragments of TRICH, or antibodies specific for TRICH may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhaner et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuvayssi, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000)

Toxicol. Lett. 112:113-67-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/news/toxicity.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are

analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for TRICH to quantify the levels of TRICH expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendozze, L. G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Selthamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference

in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schem, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in *DNA Microarrays: A Practical Approach*, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding TRICH may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1

constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) *Proc. Natl. Acad. Sci. USA* 83:7353-7357.)

Fluorescent *in situ* hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Urich, et al. (1995) in Meyers, *supra*, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding TRICH on a physical

map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps.

Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) *Nature* 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, TRICH, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between TRICH and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with TRICH, or fragments thereof, and washed. Bound TRICH is then detected by methods well known in the art. Purified TRICH can also be coated directly onto plates for use in the aforementioned drug screening techniques.

Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding TRICH specifically compete with a test compound for binding TRICH. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with TRICH.

In additional embodiments, the nucleotide sequences which encode TRICH may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below including U.S. Ser. No. 60/216,547, U.S. Ser. No. 60/218,232, U.S. Ser. No. 60/220,112, and U.S. Ser. No. 60/221,839 are expressly incorporated by reference herein, are expressly incorporated by reference herein.

## EXAMPLES

### 1. Construction of cDNA Libraries

Incye cDNAs were derived from cDNA libraries described in the LIFESeq GOLD database (Incye Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIzol (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA

purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)<sup>+</sup> RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)<sup>+</sup>PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *SUPRA*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,

PBLUESCRIPT plasmid (Stratagene), PSPT01 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## 5 II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) *Anal. Biochem.* 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labystems Oy, Helsinki, Finland).

## 20 III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families.

See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or GenScan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblies to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblies were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA.

15 The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

25 Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO.33-64. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

#### 5 IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative transporters and ion channels were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbprt and gblng). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 Kb. To determine which of these Genscan predicted cDNA sequences encode transporters and ion channels, the encoded polypeptides were analyzed by querying against PFAM models for transporters and ion channels. Potential transporters and ion channels were also identified by homology to Incyte cDNA sequences that had been annotated as transporters and ion channels. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbprt public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence.

Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbprt public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

#### "Stretched" Sequences

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or Genscan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

#### V1. Chromosomal Mapping of TRICH Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO.33-64 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO.33-64 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Patrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Genethon were used to determine if any of the clustered sequences

had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Génethon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap '99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

#### VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum (length(Seq. 1), length(Seq. 2))}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For

example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding TRICH are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding TRICH. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

#### VIII. Extension of TRICH Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg<sup>2+</sup>, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme

(Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispersing 100 µl PICOGREEN quantitation reagent (0.25% (w/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (LabSystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a 1% agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviII chloera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/Zx carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### IX. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:33-64 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

#### X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschwieler, *supra*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), *supra*). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645; Marshall, A. and J. Hodgson (1998) *Nat. Biotechnol.* 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection.

After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37° C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85° C to the stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

#### Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110° C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μl of the array element DNA, at an average concentration of 100 ng/μl, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60° C followed by washes in 0.2% SDS and distilled water as before.

#### Hybridization

Hybridization reactions contain 9 μl of sample mixture consisting of 0.2 μg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65° C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverlip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60° C. The arrays are washed for 10 min at 45° C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45° C in a second wash buffer (0.1X SSC), and dried.

#### Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate

filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

- 5 The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

- 10 The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

- 20 A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### XI. Complementary Polynucleotides

- 25 Sequences complementary to the TRICH-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring TRICH. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of TRICH. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the TRICH-encoding transcript.

#### XII. Expression of TRICH

Expression and purification of TRICH is achieved using bacterial or virus-based expression systems. For expression of TRICH in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*lac*) hybrid

- 5 promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express TRICH upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of TRICH in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographa californica nuclear polyhedrosis virus (ACMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding TRICH by either homologous recombination or bacterial-mediated

- 10 transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect *Spodoptera frugiperda* (SF9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945)

- 15 In most expression systems, TRICH is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from *Schistosoma japonicum*, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from TRICH at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoadfinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995), supra, ch. 10 and 16). Purified TRICH obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, and XVIII where applicable.

#### XIII. Functional Assays

- 30 TRICH function is assessed by expressing the sequences encoding TRICH at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which

contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of TRICH on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding TRICH and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding TRICH and other genes of interest can be analyzed by northern analysis or microarray techniques.

#### XIV. Production of TRICH Specific Antibodies

TRICH substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the TRICH amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, SUPRA, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, SUPRA.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide and anti-TRICH activity by, for example, binding the peptide or TRICH to a substrate, blocking with 1% BSA, reacting with rabbit antiserum, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

#### XV. Purification of Naturally Occurring TRICH Using Specific Antibodies

Naturally occurring or recombinant TRICH is substantially purified by immunoaffinity chromatography using antibodies specific for TRICH. An immunoaffinity column is constructed by covalently coupling anti-TRICH antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing TRICH are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of TRICH (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/TRICH binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and TRICH is collected.

#### XVI. Identification of Molecules Which Interact with TRICH

Molecules which interact with TRICH may include transporter substrates, agonists or antagonists, modulatory proteins such as G $\beta$  proteins (Reimann, SUPRA) or proteins involved in TRICH localization or clustering such as MAGUKs (Craven, SUPRA). TRICH, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled TRICH, washed, and any wells with labeled TRICH complex are assayed. Data obtained using different concentrations of TRICH are used to calculate values for the number, affinity, and association of TRICH with the candidate molecules.

Alternatively, proteins that interact with TRICH are isolated using the yeast 2-hybrid system (Fields, S. and O. Song (1989) *Nature* 340:245-246). TRICH, or fragments thereof, are expressed as fusion proteins with the DNA binding domain of Gal4 or LexA, and potential interacting proteins are expressed as fusion proteins with an activation domain. Interactions between the TRICH fusion protein and the TRICH interacting proteins (fusion proteins with an activation domain) reconstitute a transactivation function that is observed by expression of a reporter gene. Yeast 2-hybrid systems are

commercially available, and methods for use of the yeast 2-hybrid system with ion channel proteins are discussed in Niehhammer, M. and M. Sheng (1998, *Mech. Enzymol.* 293:104-122).

TRICH may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Potential TRICH agonists or antagonists may be tested for activation or inhibition of TRICH ion channel activity using the assays described in section XVIII.

#### XVII. Demonstration of TRICH Activity

Ion channel activity of TRICH is demonstrated using an electrophysiological assay for ion conductance. TRICH can be expressed by transforming a mammalian cell line such as COS7, HeLa or CHO with a eukaryotic expression vector encoding TRICH. Eukaryotic expression vectors are commercially available, and the techniques to introduce them into cells are well known to those skilled in the art. A second plasmid which expresses any one of a number of marker genes, such as  $\beta$ -galactosidase, is co-transformed into the cells to allow rapid identification of those cells which have taken up and expressed the foreign DNA. The cells are incubated for 48-72 hours after transformation under conditions appropriate for the cell line to allow expression and accumulation of TRICH and  $\beta$ -galactosidase.

Transformed cells expressing  $\beta$ -galactosidase are stained blue when a suitable colorimetric substrate is added to the culture media under conditions that are well known in the art. Stained cells are tested for differences in membrane conductance by electrophysiological techniques that are well known in the art. Untransformed cells, and/or cells transformed with either vector sequences alone or  $\beta$ -galactosidase sequences alone, are used as controls and tested in parallel. Cells expressing TRICH will have higher anion or cation conductance relative to control cells. The contribution of TRICH to conductance can be confirmed by incubating the cells using antibodies specific for TRICH. The antibodies will bind to the extracellular side of TRICH, thereby blocking the pore in the ion channel, and the associated conductance.

Alternatively, ion channel activity of TRICH is measured as current flow across a TRICH-containing *Xenopus laevis* oocyte membrane using the two-electrode voltage-clamp technique (Sisi et al., *Supra*; Jegla, T. and L. Salkoff (1997) *J. Neurosci.* 17:32-44). TRICH is subcloned into an appropriate *Xenopus* oocyte expression vector, such as pBF, and 0.5-5 ng of mRNA is injected into mature stage IV oocytes. Injected oocytes are incubated at 18°C for 1-5 days. Inside-out macropatches are excised into an intracellular solution containing 116 mM K-glucuronate, 4 mM KCl, and 10 mM Hepes (pH 7.2). The intracellular solution is supplemented with varying concentrations of the TRICH mediator, such as cAMP, cGMP, or  $\text{Ca}^{2+}$  (in the form of  $\text{CaCl}_2$ ), where appropriate.

Electrode resistance is set at 2-5 M $\Omega$  and electrodes are filled with the intracellular solution lacking mediator. Experiments are performed at room temperature from a holding potential of 0 mV. Voltage ramps (2.5 s) from -100 to 100 mV are acquired at a sampling frequency of 500 Hz. Current measured is proportional to the activity of TRICH in the assay.

In particular, the activities of TRICH-1, TRICH-2, and TRICH-10, are measured as  $\text{K}^+$  conductance, the activities of TRICH-6 and TRICH-9 are measured as  $\text{K}^+$  conductance in the presence of membrane stretch or free fatty acids, the activities of TRICH-18, TRICH-25 and TRICH-31 are measured as voltage-gated  $\text{K}^+$  conductance, TRICH-5 activity is measured as  $\text{Cl}^-$  conductance in the presence of GABA, TRICH-11 activity is measured as cation conductance in the presence of heat, and the activity of TRICH-9, TRICH-28 is measured as  $\text{Ca}^{2+}$  conductance.

Transport activity of TRICH is assayed by measuring uptake of labeled substrates into *Xenopus laevis* oocytes. Oocytes at stages V and VI are injected with TRICH mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM  $\text{CaCl}_2$ , 1mM  $\text{MgCl}_2$ , 1mM  $\text{Na}_2\text{HPO}_4$ , 5 mM Hepes, 3.8 mM NaOH, 50 $\mu\text{g/ml}$  gentamycin, pH 7.8) to allow expression of TRICH. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM  $\text{CaCl}_2$ , 1mM  $\text{MgCl}_2$ , 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids, sugars, drugs, ions, and neurotransmitters) is initiated by adding labeled substrate (e.g. radiolabeled with  $^3\text{H}$ , fluorescently labeled with rhodamine, etc.) to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in  $\text{Na}^+$ -free medium, measuring the incorporated label, and comparing with controls. TRICH activity is proportional to the level of internalized labeled substrate. In particular, test substrates include pigment precursors and related molecules for TRICH-3, aminophospholipids for TRICH-4, fructose and glucose for TRICH-7 and TRICH-15, amino acids for TRICH-8,  $\text{Na}^+$  and iodide for TRICH-12,  $\text{Na}^+$  and  $\text{H}^+$  for TRICH-13 and TRICH-21,  $\text{Na}^+$  and glucose for TRICH-16 and TRICH-19, and glucose for TRICH-23, TRICH-26, TRICH-29, TRICH-30, and TRICH-32.

ATPase activity associated with TRICH can be measured by hydrolysis of radiolabeled ATP- $[\gamma\text{-}^{32}\text{P}]$ , separation of the hydrolysis products by chromatographic methods, and quantitation of the recovered  $^{32}\text{P}$  using a scintillation counter. The reaction mixture contains ATP- $[\gamma\text{-}^{32}\text{P}]$  and varying amounts of TRICH in a suitable buffer incubated at 37°C for a suitable period of time. The reaction is terminated by acid precipitation with trichloroacetic acid and then neutralized with base, and an aliquot of the reaction mixture is subjected to membrane or filter paper-based chromatography to separate the reaction products. The amount of  $^{32}\text{P}$  liberated is counted in a scintillation counter. The amount of radioactivity recovered is proportional to the ATPase activity of TRICH in the assay.

#### XVIII. Identification of TRICH Agonists and Antagonists

TRICH is expressed in a eukaryotic cell line such as CHO (Chinese Hamster Ovary) or HEK (Human Embryonic Kidney) 293. Ion channel activity of the transformed cells is measured in the presence and absence of candidate agonists or antagonists. Ion channel activity is assayed using patch clamp methods well known in the art or as described in Example XVII. Alternatively, ion channel activity is assayed using fluorescent techniques that measure ion flux across the cell membrane (Velicelebi, G. et al. (1999) *Meth. Enzymol.* 294:20-47; West, M.R. and C.R. Molloy (1996) *Anal. Biochem.* 241:51-58). These assays may be adapted for high-throughput screening using microplates. Changes in internal ion concentration are measured using fluorescent dyes such as the  $Ca^{2+}$  indicator Fluo-4 AM, sodium-sensitive dyes such as SBFI and sodium green, or the Cl<sup>-</sup> indicator MQAE (all available from Molecular Probes) in combination with the FLIPR fluorimetric plate reading system (Molecular Devices). In a more generic version of this assay, changes in membrane potential caused by ionic flux across the plasma membrane are measured using oxonol dyes such as DiBAC<sub>4</sub> (Molecular Probes). DiBAC<sub>4</sub> equilibrates between the extracellular solution and cellular sites according to the cellular membrane potential. The dye's fluorescence intensity is 20-fold greater when bound to hydrophobic intracellular sites, allowing detection of DiBAC<sub>4</sub> entry into the cell (Gonzalez, J.E. and P.A. Negulescu (1998) *Curr. Opin. Biotechnol.* 9:624-631). Candidate agonists or antagonists may be selected from known ion channel agonists or antagonists, peptide libraries, or combinatorial chemical libraries.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Incyte	Project ID	3474637	1	3474673CDB1	33	3474673CDB1	Incyte
Polypeptide	Seq ID NO:	4588877	2	4588877CDB1	34	4588877CDB1	Polynucleotide ID
Incyte	Seq ID NO:	7472214	3	7472214CDB1	35	7472214CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7473053	4	7473053CDB1	36	7473053CDB1	Incyte
Incyte	Seq ID NO:	7473347	5	7473347CDB1	37	7473347CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7474240	6	7474240CDB1	38	7474240CDB1	Incyte
Incyte	Seq ID NO:	7475338	7	7475338CDB1	39	7475338CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7476747	8	7476747CDB1	40	7476747CDB1	Incyte
Incyte	Seq ID NO:	7477898	9	7477898CDB1	41	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	10	7477898CDB1	42	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	11	7477898CDB1	43	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	12	7477898CDB1	44	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	13	7477898CDB1	45	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	14	7477898CDB1	46	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	15	7477898CDB1	47	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	16	7477898CDB1	48	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	17	7477898CDB1	49	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	18	7477898CDB1	50	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	19	7477898CDB1	51	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	20	7477898CDB1	52	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	21	7477898CDB1	53	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	22	7477898CDB1	54	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	23	7477898CDB1	55	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	24	7477898CDB1	56	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	25	7477898CDB1	57	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	26	7477898CDB1	58	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	27	7477898CDB1	59	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	28	7477898CDB1	60	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	29	7477898CDB1	61	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	30	7477898CDB1	62	7477898CDB1	Incyte
Incyte	Seq ID NO:	7477898	31	7477898CDB1	63	7477898CDB1	Polynucleotide
Polypeptide	Seq ID NO:	7477898	32	7477898CDB1	64	7477898CDB1	Incyte

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	3474673CD1	g13507377	1.00E-151	[fl][Homo sapiens] potassium channel TASK-4 (Decher, N. et al. (2001) FEBS Lett. 492 (1-2), 84-89)
2	4588877CD1	g13926111	3.00E-96	[fl][Homo sapiens] (AF358910) 2P domain potassium channel Talk-2
3	7472214CD1	g1107730	1.70E-243	[Mus musculus] ABC8 (Savary, S. et al. (1996) Mamm. Genome 7 (9), 673-676)
		g11342541	0	[fl][Homo sapiens] putative white family ATP-binding cassette transporter
4	7473053CD1	g3850108	9.00E-209	[Schizosaccharomyces pombe] putative calcium- transporting atpase
		g3628757	0	[Homo sapiens] FIC1 (Bull, L.N. et al. (1998) Nat. Genet. 18 (3), 219-224)
5	7473347CD1	g1060975	1.70E-206	[Rattus norvegicus] GABA receptor rho-3 subunit precursor (Ogurus, T. et al. (1996) Biochim. Biophys. Acta 1305 (1-2), 15-18)
6	7474240CD1	g2745727	0	[Rattus norvegicus] potassium channel (Shi, W. et al. (1997) J. Neurosci. 17 (24), 9423-9432)
7	7475338CD1	g183298	2.10E-158	[Homo sapiens] GLUT5 protein (Kayano, T. et al. (1990) J. Biol. Chem. 265 (22), 13276-13282)
9	7477898CD1	g2745729	0	[Rattus norvegicus] potassium channel (Shi, W. et al. (1997) J. Neurosci. 17 (24), 9423-9432)
10	7472728CD1	g8452900	3.50E-261	[Rattus norvegicus] potassium channel TREK-2 (Bang, H. et al. (2000) J. Biol. Chem. 275 (23), 17412- 17419)
11	7474322CD1	g12003146	0	[fl][Homo sapiens] capsaicin receptor
12	5455621CD1	g1399954	3.00E-143	[Rattus norvegicus] thyroid sodium/iodide symporter NIS (Dai, G. et al. (1996) Nature 379 (6564), 458-460)
13	7477248CD1	g2944233	3.10E-195	[Homo sapiens] sodium-hydrogen exchanger 6 (Numata, M. et al. (1998) J. Biol. Chem. 273 (12), 6951- 6959)
14	2944004CD1	g3451312	1.40E-188	[Schizosaccharomyces pombe] membrane atpase
15	3046849CD1	g12802047	0	[fl][Homo sapiens] (AJ271290) facilitative glucose transporter GLUT11

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
16	4538363CD1	g338055	7.40E-181	[Homo sapiens] Na <sup>+</sup> /glucose cotransporter (Hediger, M.A. et al. (1989) Proc. Natl. Acad. Sci. U.S.A. 86 (15), 5748-5752)
17	6427460CD1	g6457274	0	[Mus musculus] putative E1-E2 ATPase (Halleck, M.S. et al. (1999) Physiol. Genomics (Online) 1 (3), 139-150)
18	7474127CD1	g206044	0	[Rattus norvegicus] potassium channel Kv3.2b (Wiedmann, R. et al. (1991) FEBS Lett. 288, 163-167)
19	7476949CD1	g9588428	0	[5' incm][Homo sapiens] dJ1024N4.1 (novel Sodium:solute symporter family member similar to SLC5A1 (SGLT1))
		g338055	3.70E-202	[Homo sapiens] Na <sup>+</sup> /glucose cotransporter (Hediger, M.A. et al. (1989) Proc. Natl. Acad. Sci. U.S.A. 86 (15), 5748-5752)
20	7477249CD1	g7715417	0	[Oryctolagus cuniculus] RING-finger binding protein (Mansharamani, M. et al. (2001) J. Biol. Chem. 276 (5), 3641-3649)
21	7477720CD1	g205709	0	[Rattus norvegicus] sodium-hydrogen exchange protein- isoform 4 (Orlowski, J. et al. (1992) J. Biol. Chem. 267, 9331- 9339)
22	7477852CD1	g8920219	0	[fl][Homo sapiens] epithelial calcium channel (Muller, D. et al. (2000) Genomics 67 (1), 48-53)
23	1471717CD1	g529590	5.00E-36	[Rattus norvegicus] liver-specific transport protein (Simonsen, G.D. et al. (1994) J. Cell. Sci. 107, 1065- 1072)
24	3874406CD1	g1514530	1.90E-117	[Homo sapiens] ABC-C transporter (Klugbauer, N. et al. (1996) FEBS Lett. 391 (1-2), 61- 65)
25	4599654CD1	g3242244	0	[Mus musculus] hyperpolarization-activated cation channel, HACC3 (Ludwig, A. et al. (1998) Nature 393 (6685), 587-591)

Table 3

Inyolte	3474673CD1	332	S201 S207 S234	N65 N94	Transmembrane domains: R130-M155, V245-L264 TASK R+ channel domain: V14-S332 Transmembrane domain: V139-L158 CHANNEL PROTEIN IONIC POTASSIUM SUBUNIT K+ PUTATIVE SUBFAMILY K MEMBER PD021430; A78-L162	Transmembrane domains: S430-M450, W564-D589, M618-V637 ABC transporter domain: R95-G277 ABC transporters family signature: BR00211; I100-P111, L201-D232 ABC transporters family signature: V181-D232 PROTEIN TRANSMEMBRANE TRANSPORT ATPEINING TRANSPORTER MEMBRANE ABC GLYCOPROTEIN INNER PUTATIVE PD000633; T365-Y583 do WHITE; FRUIT; FLY; SCARLET; DM05200 P45844 T89-650; G277-L623 ABC TRANSPORTERS FAMILY DM00008 P45844 73-287; I61-Q276 ABC transporter motif: L201-L215 ATP/GTP binding site (P-loop): G102-S109	HMME	HMME_PFAH	HMME	BLAST_PRODOM	HMME	BLAST_PRODOM	PROFILINSCAN	BLAST_PRODOM	BLAST_PRODOM	BLAST_DOMO	BLAST_DOMO	MOTIFS	MOTIFS
---------	------------	-----	----------------	---------	---	---	------	-----------	------	--------------	------	--------------	--------------	--------------	--------------	------------	------------	--------	--------

Table 2 (cont.)

polypeptide seq ID NO:	Incyte polypeptide ID	GenBank ID No:	GenBank ID Probability score	GenBank Homolog
26	5047435CD1	G13445575	0	[f1][Homo sapiens] facilitative glucose transporter GLUT10 (McVie-Wylie, A.J. et al. (2001) Genomics 72 (1), 113-117)
27	7475603CD1	G9211112	0	[f1][Homo sapiens] macrophage ABC transporter (Kaminski, W.E. et al. (2000) Biochem. Biophys. Commun. 273 (2), 532-538)
28	7477845CD1	G3800830	0	[Rattus norvegicus] putative four repeat ion channel (Lee, J.H. et al. (1999) FEBS Lett. 445 (2-3), 231-236)
29	168827CD1	G7707622	1.20E-116	[Homo sapiens] organic anion transporter 4 (Chen, S.H. et al. (2000) J. Biol. Chem. 275 (6), 4507-4512)
30	7472734CD1	G7707622	4.50E-117	[f1][Rattus norvegicus] putative integral membrane transport protein (Schomberg, E. et al. (1998) FEBS Lett. 425 (1), 79-86)
31	7473473CD1	G6625694	0	[Rattus norvegicus] potassium channel Eag2 (Sagami, M.J. et al. (1999) J. Neurosci. 19 (24), 10789-10802)
32	7477725CD1	G3004482	1.00E-177	[f1][Rattus norvegicus] putative integral membrane transport protein (Schomberg, E. et al. (1998) FEBS Lett. 425 (1), 79-86)
	G7707622	4.20E-130		[Homo sapiens] organic anion transporter 4 (Chen, S.H. et al. (2000) J. Biol. Chem. 275 (6), 4507-4512)

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4	7473053CD1	1190	S153 S259 S268 S391 S413 S452 S493 S545 S573 S624 S631 S687 S723 S739 S744 S832 S1174 S1132 S1164 S1124 S1143 S1168 T267 T36 T370 T378 T514 T519 T580 T646 T705 T732 T899 T980 T1098 T1158 Y23 Y29 Y489 Y607	N579	Transmembrane domains: S77-V94, L276-W298, Y330-R350, L947-I971, Q991-I1009 E1-E2 ATPase domains: E381-V403, Q530-A562, Y633-G685, R788-D818 E1-E2 ATPases phosphorylation site proteins BL00154: G134-L151, V386-F404, D650-M690, T809-S832 E1-E2 ATPases phosphorylation site: A372-V417 P-type cation-transporting ATPase superfamily signature PR00119: F390-F404, A666-D676, I812-I831 ATPASE HYDROLASE TRANSMEMBRANE PHOSPHORYLATION ATPBINDING PROTEIN PROBABLE CALCIUMTRANSPORTING CALCIUM TRANSPORT PD004657: S846-F1093 FIC1 PROTEIN PD180313: H1039-W1165 do ATPASE; CALCIUM; TRANSPORTING; DM02405 P32660 318-1225: W128-F418, E466-N910 ATPase E1-E2 motif: D392-T398 Transmembrane domain: V332-V351	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLIMPS_PRINTS BLAST_PRODROM BLAST_PRODROM BLAST_DOMO MOTIFS
5	7473347CD1	467	S149 S175 S344 S37 S390 S411 S419 S427 S53 S96 T100 T136 T157 T355 T356 T366 T41	N126 N197 N220	Transmembrane domain: V332-V351	HMMER

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
5					Neurotransmitter-gated ion-channel domain: P58-Q362, H441-W463 Neurotransmitter-gated ion channels signature BL00236: V85-P122, I139-H148, D169-Y207, Y254-A295 Neurotransmitter-gated ion-channels signature: L164-H218 Neurotransmitter-gated ion-channels signature PR00252: T105-F121, K138-S149, C184-C198, S261-P273 Gamma-aminobutyric acid A (GABAA) receptor signature PR00253: F270-W290, V296-V317, V330-V351, Y446-Y466 CHANNEL IONIC TRANSMEMBRANE GLYCOPROTEIN POSTSYNAPTIC MEMBRANE RECEPTOR PRECURSOR SIGNAL PROTEIN PD000153: E62-S427 NEUROTRANSMITTER-GATED ION-CHANNELS DM00560 P50573 34-464: S37-V467 Neurotransmitter-gated ion channels motif: C184-C198	HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLIMPS_PRINTS BLIMPS_PRINTS BLAST_PRODROM BLAST_DOMO MOTIFS

Seq ID	Incyte ID	Polypeptide	Amino Acid	Phosphorylation	Glycosylation	Potential	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7							SUGAR TRANSPORT PROTEINS DM00135 P22732 132-466; R138-T473 SUGAR transporter 1 motif: S338-A53 SUGAR transporter 2 motif: V140-R165 HMMEP HMMEP_PFAAM TRANSMEMBRANE AMINO ACID TRANSPORTER A102-G543 PROLINE PD001875: W80-L380 TRANSMEMBRANE DOMAIN: L300-N318 CYCLIC NUCLEOTIDE-BINDING DOMAIN: V608-A699 POTASSIUM CHANNEL IONIC CHANNEL PD118772: E702-S955 CHANNEL PROTEIN IONIC POTASSIUM NONPHOTOPHOBIC HYPOCOTYL PUTATIVE SUBUNIT REPEAT EAG PD009483: M1-L86 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 I38465 562-948; H413-P738 do POTASSIUM CHANNEL; KST1; AKT1; DM02383 I38465 353-560; T201-A412 BLAST_DOMO	BLAST_DOMO
8	7476747CD1		S143 S365 S4 S456 S46 S51 S55 T34 T430 Y45 N141 N205 N214 N256 N562 N62 N76				TRANSMEMBRANE DOMAINS: 1242-P269, Y289-P308, I322-Y342 HMMEP HMMEP_PFAAM TRANSMEMBRANE AMINO ACID TRANSPORTER A102-G543 ACID AMINO PROTEIN TRANSPORTER PERMEASE TRANSMEMBRANE INTERGENIC REGION PUTATIVE PD001875: W80-L380 TRANSMEMBRANE DOMAIN: L300-N318 CYCLIC NUCLEOTIDE-BINDING DOMAIN: V608-A699 POTASSIUM CHANNEL IONIC CHANNEL PD118772: E702-S955 CHANNEL PROTEIN IONIC POTASSIUM NONPHOTOPHOBIC HYPOCOTYL PUTATIVE SUBUNIT REPEAT EAG PD009483: M1-L86 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 I38465 562-948; H413-P738 do POTASSIUM CHANNEL; KST1; AKT1; DM02383 I38465 353-560; T201-A412 BLAST_DOMO	BLAST_DOMO
9	7477898CD1		S105 S140 S145 S200 S26 S283 S288 S458 S488 S55 S670 S706 S724 S751 S774 S788 S864 S872 S879 S897 S929 T13 T170 T202 T220 T301 T326 T363 T377 T486 T522 T678				TRANSMEMBRANE DOMAIN: L300-N318 CYCLIC NUCLEOTIDE-BINDING DOMAIN: V608-A699 POTASSIUM CHANNEL IONIC CHANNEL PD118772: E702-S955 CHANNEL PROTEIN IONIC POTASSIUM NONPHOTOPHOBIC HYPOCOTYL PUTATIVE SUBUNIT REPEAT EAG PD009483: M1-L86 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 I38465 562-948; H413-P738 do POTASSIUM CHANNEL; KST1; AKT1; DM02383 I38465 353-560; T201-A412 BLAST_DOMO	BLAST_DOMO

Table 3 (cont.)

Seq ID	Incyte ID	Polypeptide	Amino Acid	Phosphorylation	Glycosylation	Potential	Signature Sequences, Domains and Motifs	Analytical Methods and Databases	
6	7474240CD1		S174 S187 S209 S211 S239 S269 S274 S275 S317 S349 S354 S514 S55 S609 S639 S821 S869 S879 S883 S896 S899 S906 S922 S923 S939 S940 S963 S974 S985 S1020 S1091 S1170 S1096 T133 T169 T344 T371 T392 T528 T582 T637 T673 T74 T829 T857 T916 T1022 T1027 T1134 T1099 Y248 Y446 Y98					TRANSMEMBRANE REGION CYCLIC NUCLEOTIDE-BINDING DOMAIN: Y492-T171 GATED ION CHANNEL: TRANSMEMBRANE REGION CYCLIC NUCLEOTIDE-BINDING DOMAIN: Y551-Y571 TRANSMEMBRANE DOMAIN: HMMEP HMMEP_PFAAM CYCLIC NUCLEOTIDE-BINDING DOMAIN: M759-E850 POTASSIUM CHANNEL IONIC CHANNEL PD104127: S852-Y1028 POTASSIUM CHANNEL IONIC CHANNEL PD104126: A1076-K1196 BLAST_PRODOM BLAST_PRODOM CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 I38465 562-948: H566-A914 do POTASSIUM CHANNEL; KST1; AKT1; DM02383 I38465 353-560: S353-A563 do CHANNEL; POTASSIUM; EAG; DM05484 I38465 1-351: M1-P351 Signal peptide: M1-A35 Transmembrane domains: C79-G96, M171-L188, Y322-V342, P448-I466 Sugar (and other) transporter domain: A26-P481 Sugar transport proteins signatures: A119-I185, V323-S379 Sugar transporter signature PR00171: A35-V45, V135-M154, Q294-Y304, I383-V404, T406-P418 Glucose transporter signature PR00172: L284-Y305, Q321-V342, L352-Q372, I383-T406, A416-P434, Y446-I466 BLIMPS-PRINTS BLIMPS-PRINTS	BLAST_DOMO BLAST_DOMO BLAST_PRODOM BLAST_PROD

Table 3 (cont.)

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
10	7472728CD1	724	S229 S283 S303 S333 S512 S545 S597 S666 S718 T104 T19 T223 T444 T515 T540 T557 T591 T636 T640 T650 T661 T676	N327 N330 N331 N532 N664 N684 N716	Transmembrane domains: A370-L388, I419-F437, V486-M503 TASK K+ channel domain: M250-D646 TWIK1 RELATED POTASSIUM CHANNEL, SUBFAMILY K, MEMBER 2 TREK1 K+ CHANNEL SUBUNIT IONIC CHANNEL PD085853: P215-G326	HMNER HMNER_PFBM BLAST_PRODUM
11	7474322CD1	470	S134 S142 S245 S326 S355 S408 S411 S415 S432 S452 T15 T22 T229 T265 T337 T341 T36	N236 N256 N321 N380	Transmembrane domains: P62-Y87, F139-F163, F212-L230, I293-I312 VANILLOID RECEPTOR SUBTYPE 1 PD137334: C348-K470	HMNER BLAST_PRODUM
12	5455621CD1	618	S110 S265 S313 S373 S490 S550 S565 S576 S594 T154 T237 T268 T360 T37 T526 T567 T70	N219 N256 N480 N574	Transmembrane domains: D10-F28, F81-Y104, F278-M297, L439-Y459, I502-R528 Sodium:solute symporter family domain: F41-G445 Sodium:solute symporter signature BL00456: T154-G208 Sodium:solute symporter family signature: N151-T198 TRANSMEMBRANE TRANSPORT PERMEASE PROTEIN SODIUM SYMPORTER PROLINE COTRANSPORTER SYMPORTER GLYCOPROTEIN PD000991: F41-C304 SYMPORTER SODIUM IODIDE THYROID SODIUM/IODIDE NIS PD024705: I446-L489, S490-G575 SODIUM:SOLUTE SYMPORTER FAMILY DM00745 P31636 24-561: D10-N219, G220-Y459	HMNER HMNER_PFBM BLIMPS_BLOCKS PROFILES SCAN BLAST_PRODUM BLAST_PRODUM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
13	7477248CD1	631	S149 S212 S258 S522 S9 T518 T551 T73 T79 Y14	N352 N516 N96	Transmembrane domains: V22-F41, L159-M181, I391-A407 Sodium/hydrogen exchanger family domain: L25-V491 Na+/H+ exchanger isoform 6 signature PR01088: Y14-I38, W39-V57, Y58-V84, Q119-E132, A269-M288, T480-Q506, K515-D533, P539-Q567, P566-E593 Na+/H+ exchanger signature PR01084: I133-F144, G147-S161, I162-T170, G208-T218 + TRANSPORT EXCHANGER NA PD01672: I133-M181 NA+/H+ PROTEIN TRANSMEMBRANE TRANSPORT ANTI PORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: G20-G63, E132-R490 SODIUMHYDROGEN EXCHANGER 6 MYELOBLAST KIAA0267 PD177855: G478-Y591 do BETA; EXCHANGER; NA; DM02572 P48764 10-734: L124-L541	HMNER HMNER_PFBM BLIMPS_PRINTS BLIMPS_PRINTS BLIMPS_PRODUM BLAST_PRODUM BLAST_PRODUM BLAST_DOMO
14	2944004CD1	1256	S103 S130 S144 S170 S227 S252 S523 S802 S817 S899 S901 S98 S1055 T269 T353 T358 T387 T502 T549 T576 T74 T912 T1212 T1061 T1236 Y349 Y407	N150 N23 N300 N312 N318 N704 N1045 N1053 N1059 N1073 N1247	Transmembrane domains: Y231-Y251, L415-L434, I933-I959, F966-L985, I1002-F1020, N1104-M1122 E1-E2 ATPase domains: V274-V365, G490-D506, Q672-A785, L851-S899 E1-E2 ATPases phosphorylation site signature BL00154: V454-G490, L492-L510, K652-C662, N724-M764, V878-S901, A905-V938 E1-E2 ATPases phosphorylation site: I478-E526	HMNER HMNER_PFBM BLIMPS_BLOCKS PROFILES SCAN

Table 3 (cont.)

530	Incyte	4538363CD1	596	potential	phosphorylation	S17 S290 S39 S5	1119 7211	N4 N545	N239 N386	N96	Transmembrane domains: S73-W95, I185-I212, L356-A376, L410- V430, F473-F491, Y513-L533	HMHER	HMHER	HMHER_PAM	Sodium:solute symporter family domain: Y50-G479	Sodium:solute symporter signature BL00456: Y27-G81, A103-R132, L165- G219, P452-G461	Sodium:solute symporter family signatures: H162-I209, V412-D502	TRANSMEMBRANE TRANSPORT PERMEASE SODIUM SYMPORT PROLINE COTRANSPORTER SYMPORTER GLYCOPROTEIN PD000991: Y50-G479	NA+/GLUCOSE COTRANSPORTER-RELATED PROTEIN PD14393: L551-A596	NA+/GLUCOSE COTRANSPORTER-RELATED PROTEIN PD166538: M1-G49	SODIUM:SOLUTE SYMPORTER FAMILY DM00745 P13866 24-561: S17-W548	BLAST_DOMO	MOTIFS	Na:solute symporter 2 motif: G461-V481
-----	--------	------------	-----	-----------	-----------------	-----------------	-----------	---------	-----------	-----	--	-------	-------	-----------	--	--	---	--	---	---	---	------------	--------	---

Table 3 (cont.)

Seq ID	Incyte ID	Amino Acid	Potential phosphorylation Sites	Potential glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
14	NO: 14	Residues				
15	3046849CD1	499	S100 S118 S215 S285 T466 T487	N292 N34 M50	<p>Transmembrane domains: M163-L181, T371-G389, M418-L440</p> <p>Sugar (and other) transporter signature: L118-L474</p> <p>Sugar transporter proteins signature: A112-V178</p> <p>Sugar transporter signature PR00171: T28-I38, M128-M147, M376- T397, T399-C411</p> <p>Glucose transporter signature PR00172: Q314-I335, M376-T399, A409- L427</p> <p>SUGAR TRANSPORT PROTEINS DM00135 P22732 I32-466: R131-T466</p> <p>Sugar transporter Z motif: T133-R158</p>	<p>BLAST_PRINTS</p> <p>BLAST_PRODOM</p> <p>BLAST_DOMO</p> <p>MOTIFS</p> <p>SPSCAN</p> <p>HMMER</p> <p>HMMER_PFAH</p> <p>PROFILERSCAN</p> <p>BLAST_PRINTS</p> <p>BLIMPS_PRINTS</p>

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
17	6427460CD1	1192	S143 S169 S188 S283 S287 S335 S451 S507 S508 S52 S555 S561 S722 S933 T203 T255 T259 T269 T333 T380 T413 T418 T659 T708 T714 T715 T910 T1103 T1017 T1105 Y885 Y1026	N397 N745 N921 N989 N1001	Transmembrane domains: V299-Y316, F1004-L1022, I1030-W1049, A1075-L1092 E1-E2 ATPase domains: E403-E425 I550-C698 E1-E2 ATPases phosphorylation site signature BL00154: G149-F166, V408-F426, D563- L703 E1-E2 ATPases phosphorylation site: L395-C442 P-type cation-transporting ATPase superfamily signature PR00119: F412-F426, A679-D689 ATPASE HYDROLASE TRANSMEMBRANE PHOSPHORYLATION ATPBINDING PROTEIN PROBABLE CALCIUMTRANSPORTING CALCIUM TRANSPORT PD004657: A857-V1108 do ATPASE; CALCIUM; TRANSPORTING; DM02405 Q09891 206-1107: T105-Y436, F471-N921 E1-E2 ATPase motif: D414-T420	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLIMPS_PRINTS BLAST_PRODROM BLAST_DOMO MOTIFS
18	7474127CD1	638	S205 S224 S336 S378 S414 S541 S553 S564 S86 T120 T146 T155 T17 T21 T25 T283 T374 T49 T520 T546 T579	N259 N266 N518 N536 N84	Transmembrane domains: I231-L248, F382-Y401, M451-V473 Ion transport protein domain: L240-I472 Potassium channel signature PR00169: E101-T120, P222-T250, Y284- K307, F310-V330, F352-S378, E381-E404, F421-M443, G450-F476	HMMER HMMER_PFAM BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
18					VOLTAGEGATED POTASSIUM CHANNEL PROTEIN KV3.2 KSHIIIA IONIC TRANSMEMBRANE ION TRANSPORT GLYCOPROTEIN MULTIGENE FAMILY ALTERNATIVE SPLICING PHOSPHORYLATION PD085814: K495-S538 do CHANNEL; POTASSIUM; CDRK; FORM; DM00436 P22462 189-350: R189-R351 do CHANNEL; POTASSIUM; CDRK; SHAW; DM00490 P22462 34-151: L34-C152	BLAST_PRODROM BLAST_DOMO BLAST_DOMO
19	7476949CD1	681	S307 S421 S56 S573 S582 S587 S638 S651 T422 T485 T650 Y510	N113 N251 N256 N403 N603	Transmembrane domains: I38-I57, S90-W112, I150-I167, L188- M207, L373-A393, V432-I448, Y530-L550 Sodium:solute symporter family domain: Y67-G496 Sodium:solute symporter signature BL00456: Y44-G98, A120-R149, L182- G236, P469-A478 Sodium:solute symporter family signatures: Q179-V226, D458-D519 TRANSMEMBRANE TRANSPORT PERMEASE PROTEIN SODIUM SYMPORT PROLINE COTRANSPORTER SYMPORTER GLYCOPROTEIN PD000991: Y67-G496 SODIUM:SOLUTE SYMPORTER FAMILY DM00745 P13866 24-561: H34-W565 Na solute symporter 1 motif: G183-A208	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILES SCAN BLAST_PRODROM BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyste Polypeptide	Amino Acid	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
21					Na <sup>+</sup> /H <sup>+</sup> exchanger isoform 2 (NHE2) signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_DOMO HMMER_Pfam HMMER MOTIFS BLAST_PRODOM HMMER_Pfam HMMER BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases
22					Signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases
23					Signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases

Table 3 (cont.)

SEQ ID NO:	Incyste Polypeptide	Amino Acid	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
20					Signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases
21					Signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases
22					Signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases
23					Signature P101086, P115-S128, K616-I627 + TRANSPORT EXCHANGER NA P101672, A83-I113, I129-L177, Y178-L212, A213-P249, D262-I287, S288-Y321, L322-M355, S359-P405, Y406-P452, I489-K531, I532-G562, R593-R640 NA <sup>+</sup> /H <sup>+</sup> PROTEIN TRANSMEMBRANE TRANSPORT ANTIPORTER SYMPORT SODIUM EXCHANGER GLYCOPROTEIN SODIUM/HYDROGEN PD000631: I77-A438 do BETA; EXCHANGER; NA; DM02572 P26434 I4-I16; L15-L687 Transmembrane domains: P493-P512, M554-M570 Ankyrin repeats: L178-S108, A116-T148, P162-S194 VANILLOID RECEPTOR SUBTYPE 1 PD101189: F115-L220 ATP/GTP binding site (P-loop): A412-T419 transmembrane domain: I48-V71, V86-P104, Y172-I199, I199-V217, P384-P402, V452-C472 Sugar (and other) transporter: I48-K492 SUGAR TRANSPORT PROTEINS DM00032 P30638 80-152; R45-K115 VESICLE; SYNAPTIC; SV2; FORM DM08835 S34961 180-344; I119-N249	BLAST_DOMO BLAST_PRODOM BLIMPS_PRODOM BLIMPS_PRODOM BLIMPS_PRINTS Databases

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
24	3874406CD1	1494	S30 S50 S134 S230 S368 S549 S638 S669 S686 S696 S792 S800 S831 S912 S1004 S1070 S1146 S1172 S1206 S1365 T111 T435 T449 T501 T520 T632 T649 T657 T729 T845 T1049 T1134 T1217 T1247 T1295 T1318 T1339 T1422 T1482 Y824	N109 N130 N313 N421 N453 N71 N788 N817 N84 N867 N91 N1182	transmembrane domain: L204-F221, T272-L290, L735-Y753, P896-S914, V941-I959, L975-R998, F1019-V1039 ABC transporter: G384-G566 G1190-G1366 ABC transporters family proteins BL00211: I389-L400, L492-D523 ABC transporters family signature: V472-D523 ABC TRANSPORTERS FAMILY DM00008 P41233 839-1045:I355-N565, K1177-M1363 DM00008 P34358 611-816:I355-N565, A1179-M1363 DM00008 P41233 1851-2058:K1173-S1365, I355-N565 DM00008 P23703 41-246:E1162-G1366, L377-G566 ATP/GTP-binding site motif A (P-loop): G391-S398, G1197-2004	HMMER HMMER_PFAM BLIMPS_BLOCKS PROFILESCAN BLAST_DOMO MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
25	4599654CD1	774	S355 S356 S40 S505 S552 S559 S597 S61 S67 S734 S736 T203 T418 T668 T764 Y490	N291 N416	transmembrane domain: Y95-F118, T203-L219, L327-L353 Transmembrane region cyclic Nucleotide G: Y168-I414 Cyclic nucleotide-binding domain: K443-M531 Cyclic nucleotide-binding domain proteins BL00888: G452-V475, G488-L497 cAMP-dependent protein kinase signature PR00103: F449-R463, S489-T498 HYPERPOLARIZATIONACTIVATED CATION CHANNEL, HAC3 PD180735: T538-M774 CHANNEL IONIC POTASSIUM K+ SUBUNIT HYPERPOLARIZATIONACTIVATED PROTEIN PUTATIVE EAG LONG PD001039: E74-R167 CAMP RECEPTOR PROTEIN CYCLIC NUCLEOTIDE-BINDING DOMAIN DM01165 A55251 333-706:H263-P561 DM01165 P29973 311-684:H263-P561 DM01165 Q03041 286-658:H263-G548 DM01165 S52072 262-635:H263-Q595	HMMER HMMER_PFAM HMMER_PFAM BLIMPS_BLOCKS BLIMPS_PRINTS BLAST_PRODOM BLAST_PRODOM BLAST_DOMO

[illegible]

Table 3 (cont.)

26	5047435CD1	614	S116 S210 S290 S538 S577 S606 T267 T432 T443 T591	N407 N599	transmembrane domain: V124-I144, A168-M190, A371-V390, W483- I511, S526-I543, F552-V570 Sugar (and other) transporter: L83-F585 Sugar transport proteins BL00216: L174-S223, G92-S103 Sugar transporter signature PR00171: G92-I102, V175-I194, L486- V507, S509-F521 Glucose transporter signature PR00172: V343-V364, L486-S509, R519- L537, W550-V570 Sugar_transport_1: G138-G153 A360-A375 Sugar transport proteins signatures sugar_transport_1.pr1: L344-S401 sugar_transport_2.pr1: A160-A225 SUGAR TRANSPORT PROTEINS DM00135 S25015 I22-478:A160-D417, L480-K574, DM00135 P09830 I01-452:G161-V405, L481-K574 DM00135 Q01440 I01-433:M178-G388, L486-G388, L486-G575 DM00135 P15729 I242-463:A485-S577, K286-L414	BLAST_DOMO	PROFILES_CAN	MOTIFS	BLIMPS_PRINTS	BLIMPS_PRINTS	BLIMPS_BLOCKS	HMMER_PFM	HMMER	Methods and Analytical Databases
----	------------	-----	--	-----------	---	------------	--------------	--------	---------------	---------------	---------------	-----------	-------	--

Table 3 (cont.)

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
27					ABC TRANSPORTERS FAMILY DM00008 P41233 839-1045:V841-A1046, L1829-M2032 DM00008 P41233 1851-2058:V1826-N2034, V841-V1045 DM00008 P34358 1441-1640:L1827-M2032, V843-V1045	BLAST_DOMO
28	7477845CD1	1737	S23 S254 S687 S692 S695 S7 S713 S766 S773 S8 S861 S1113 S1228 S1271 S1455 S1463 S1537 S1595 S1647 S1652 S1730 T272 T324 T886 T1257 T1320 T1359 T1387 T1406 T1456 T1486 T1528 T1561 T1570 T1645 T1694 Y419 Y702 Y832	N210 N216 N859 N1064 N1371 N1449	transmembrane domain: M1244-A1262, V1319-F1336, I1338-F1357, A1423-I1446, W107-V126, V181-M199, S298- I321, L509-V531, V575-I598, Y879-M904, I1017-F1034, I1134-V1152 Ion transport protein ion_trans: W32-I321 M380-I598 L884-V1155 I1206- I1446 Calcium channel signature PR00167: D535-D561 PROTEIN F17C8.6 C11D2.5 NEARLY IDENTICAL C ELEGANS PREDICTED PD023984: V1447-S1637, E1714-T1720 C11D2.6 PROTEIN PD178227: L1241-R1368, I1206-F1292 P585-E606 C11D2.6 PROTEIN SIMILARITY ALONG ENTIRE GENE CALCIUM CHANNEL ALPHA PROTEINS PD041964: L599-V885, CHANNEL CALCIUM IONIC SUBUNIT VOLTAGE GATED SODIUM ALPHA TRANSMEMBRANE L TYPE PD000032: Y887-V1120, I33-V330, K1361- F1450, I1206-F1357, I577-I598, F1337- L1356, I1134-F1159, D1416-V1443	HMNER  HMNER_PFAM  BLIMPS_PRINTS  BLAST_PRODOM  BLAST_PRODOM  BLAST_PRODOM  BLAST_PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
28					III REPEAT DM00079 A55138 1052-1268:V1020-L1227 DM00079 P35500 1424-1636:W1090-P1194, I1017-N1050 IV REPEAT DM00277 P27732 1363-1572:F1337-L1536 DM00277 P15381 1384-1595:F1337-L1536	BLAST_DOMO  BLAST_DOMO
29	168827CD1	547	S109 S167 S201 S282 S336 S404 S408 S526 T133 T323 T35 T432 T453 T58	N102 N107 N56	transmembrane domain: F16-T35, Y180-C200, S201-V222, M410- E429, T469-Y492, L496-L514 Sugar (and other) transporter: L13-Q528 ORGANIC TRANSPORTERLIKE TRANSPORT PROTEIN RENAL ANION TRANSPORTER CATIONIC KIDNEYSPECIFIC SOLUTE PD151320: N102-L144	HMNER  HMNER_PFAM  BLAST_PRODOM
30	7472734CD1	547	S143 S167 S201 S282 S336 S404 S408 S46 S526 S60 S68 T133 T323 T432 T453 T58	N102 N39 N56 N62	transmembrane domain: I18-F32, M147-Y163, Y180-C200, S201- V222, M410-E429, T469-Y492, L496-L514 Sugar (and other) transporter: L18-Q528 SUGAR TRANSPORT PROTEINS DM00032 P46501 280-351:V121-K173 ORGANIC TRANSPORTERLIKE TRANSPORT PROTEIN RENAL ANION TRANSPORTER CATIONIC KIDNEYSPECIFIC SOLUTE PD151320: N102-K145	HMNER  HMNER_PFAM  BLAST_DOMO  BLAST_PRODOM

[illegible]

Table 3 (cont.)

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
33	3474673CB1	1775	1-391, 578-786, 1024-1301	GNFL.g7798848_00000 3.004.edit 6724643H1 (LUNLMT01) 3474673H1 (LUNGNOT27) 71495515V1 71495515V1 FL135171_00001 71497982V1	1 861 249 1205 975 539 1	1156 1347 568 1775 1545 1534 662
34	4588877CB1	1545	261-619, 1-193, 794-1071	GBI:g8117242_000054 edit.8639-8803 GBI:g8117242_000054 edit.4857-4997 GBI:g8117242_000054 edit.10305-10463 6891360H1 (BRAITDR03) GBI:g8117242_000054 edit.50-89 GBI:g8117242_000054 edit.6950-7093 GBI:g8117242_000054 edit.4345-4478 60124962D2 GBI:g8117242_000054 edit.8313-8414 GBI:g8118985_000043 edit.12301- 12444.comp GBI:g8117242_000054 edit.4112-4228 GBI:g8117242_000054 edit.10957-11181 5500380H1 (BRABDIR01) GBI:g8117242_000054 edit.10616-10732	1171 544 1441 1433 1 925 358 1735 1069 685 241 1717 907 1600	1335 684 1599 1905 240 1068 492 1941 1170 810 357 1941 1119 1716
35	7472214CB1	1941	1483-1558, 1- 413, 495-616, 732-1149			

WO 02/04520

PCT/US01/21448

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
35				GBI:g8117242_000054 edit.8907-9011 GBI:g8117242_000054 edit.6643-6756	1336 811	1440 924
36	7473053CB1	4971	3312-3482, 1- 1466, 4307-4971, 2184-2221	8035016H1 (SMCRUNE01) 6822202J1 (SINTNOR01) 6781747H1 (OVARDIR01) 8035016J1 (SMCRUNE01) 6824230H1 (SINTNOR01) 6894266H1 (BRAITDR03) 6777836H1 (OVARDIR01) 6908503H1 (PITUDIR01) 6908503J1 (PITUDIR01) 6823447H1 (SINTNOR01) 6823447J1 (SINTNOR01) 6006310F8 (FIBRUNT02) 4171959T6 (SINTNOT21) 5088860F6 (UTRSTMR01)	2315 2145 968 2979 2867 548 1601 1 1270 3525 4226 4501 3637 4461	2975 2877 1449 3643 3483 1157 2238 667 1830 4260 4829 4969 4287 4853
37	7473347CB1	1404	126-633, 1013- 1404, 768-838	GBI.lee4.edit	1	1404

WO 02/04520

PCT/US01/21448

Table 4 (cont.)

Polynucleotide	Incyte	Polynucleotide ID	Sequence Length	Selected Sequence	Sequence Fragments	5' Position	3' Position
38		7474240CB1	4048	3023-4048, 1753-2469, 1-920, 1593-1658, 2614-2908, 1138-1367	71984804V1 GBI:g756646 edit 71986624V1 55055014H1 55037111J2 7198368V1 2612 4048 55037119J2 2502027F6 (ADPRT05)	964 929 1976 130 871 2043 2612 4048 875 1235	1311 3418 1976 130 871 2043 2612 4048 875 1235
39		7475338CB1	1539	1412-1539, 1-328, 495-837, 922-1218	GBI:g7960701.000004 edit. 549-713 GBI:g7960701.000004 edit. 13381-13480 GBI:g7960701.000004 edit. 8755-8943 GBI:g7960701.000004 edit. 4292-4417 GBI:g7960701.000004 edit. 16237-16317 GBI:g7960701.000004 edit. 16237-16317 GBI:g7960701.000004 edit. 20107-20325 GBI:g7960701.000004 edit. 9989-10099 GBI:g7960701.000004 edit. 18748-18873 GBI:g7960701.000003 edit. 9783-9884 GBI:g7960701.000004 edit. 5251-5403 GBI:g7960701.000004 edit. 8384-8506 71906448V1 71534467V1 3351512F6 (PROSNOT28) 7761783J1 (THYMNOE2) 6934981R8 (SINTMR2)	154 1015 715 438 1194 1321 1539 1014 1320 153 591 714 1082 1539 2724 2570 860	312 1113 903 438 1194 1321 1539 1014 1320 153 591 714 1082 1539 2724 2570 860
40		7476747CB1	3114	1717-1870, 1-503, 1468-1650	6934981R8 (SINTMR2)	78	860

114

Table 4 (cont.)

Polynucleotide	Incyte	Polynucleotide ID	Sequence Length	Selected Sequence	Sequence Fragments	5' Position	3' Position
40					6393368H1 (PROSNOT1) 70536163V1 6934981R8 (SINTMR2) GNN.g7712065_000012 7080657H1 (STOMTR02) 5633289H1 (PLACFER01) 85746200 GBI:g2262095	1782 2575 3114 643 1922 1403 890 1473 2877	2075 3114 643 1922 1403 890 1473 2877
41		7477898CB1	2877	846-901, 1272-1378, 2319-2877	55022826J1 55030210H1 43936676 (TESTTUT03) 55030274H1 8565876 55018149J1 FL203597_00001 GNN.g7263861_026.ed 1t	1138 403 2231 1482 2597 1907 712 1052	1834 986 2777 2153 2820 2585 1807 1052
42		7472728CB1	2820	1-1399, 2207-2229	55022826J1 55030210H1 43936676 (TESTTUT03) 55030274H1 8565876 55018149J1 FL203597_00001 GNN.g7263861_026.ed 1t	1138 403 2231 1482 2597 1907 712 1052	1834 986 2777 2153 2820 2585 1807 1052
43		7474322CB1	1440	1-604, 714-768	GBI:g8081632 edit 71228887V1 70868623V1 3696546T6 (SINTMR05) 706744954V1 1426382H1 (SINTMR05) 3696546T6 (SINTMR05) 6828352H1 (SINTMR05) 3699565H1 (SINTMR05) 7700096H1 (KIDPTD01) 70678552V1	1 1090 988 1833 1520 1224 799 530 1 250 1419	1440 1440 1385 2394 2091 1492 1381 1149 281 990 2055
44		5455621CB1	2394	1483-1686, 1-329, 838-1155, 2201-2235	70678552V1 (KIDPTD01)	1419	2055

115

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
45	7477248CB1	2890	1-58, 2739-2890, 2310-2349, 329- 1167	2777287H1 (OVARTUT03)	2250	2498
				7977733H1 (LSUBDMC01)	841	1427
				7678168J1 (NOSETUE01)	1271	1827
				7611941J1 (KIDCTME01)	2273	2890
				6590507H1 (TLYMUNT03)	179	672
				2701794F6 (OVARTUT10)	1208	1741
				2544096F6 (UTRSNOT11)	1732	2252
				60117044D2	1	431
				5020832H1 (OVARNON03)	2195	2471
				7662529H1 (UTRSTME01)	526	926
				4762728F6 (PLACNOT05)	872	1387
				g2264624	2268	2446
				6264977H1 (MCLDTXN03)	1210	1797
				2944004F6 (BRAITUT23)	2790	3531
				6610392H2 (MUSTTMC01)	3306	3926
				GNN.g7328818_000024 002.edit	2145	2648
46	2944004CB1	3926	3338-3365, 1- 687, 1222-2267	7035078H1 (SYNTFER03)	1	440
				7620248J1 (HEARFEE03)	2431	3039
				496537H1 (HNT2NOT01)	2329	2487
				6264427T8 (MCLDTXN03)	453	1174
				6264427F8 (MCLDTXN03)	170	842

WO 02/04520

PCT/US01/21448

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
46				7673654H1 (FIBPFEC01)	1733	2239
47	3046849CB1	2135	2072-2135, 596- 711, 1014-1263	8262790U1	1383	2135
				71896642V1	1	592
				71247870V1	1050	1736
				FL3046849_g6815043_ 000004.g183298	51	1520
48	4538363CB1	2637	1-183, 1575- 1680, 2094-2637	FL4538363_g3126781_ g520469	1	1917
49	6427460CB1	3783	985-1833, 2687- 3204	71401405V1	1766	2637
				70857895V1	416	1035
				7727961J1 (UTRCDE01)	3284	3783
				70857789V1	566	1109
				g5689372.edit	1092	3361
50	7474127CB1	2105	1078-2105	g3801917	1	452
				GBI.g8568959.edit_3	1119	2105
				g6140313	482	951
				5819744F7 (PROSTUS23)	168	479
				g5920552	1	488
51	7476949CB1	2069	1233-1356, 1- 117, 2047-2069, 347-503, 1536- 1844	55049678J1	862	1359
				FL7476949_g6714723_ g338053	1	2046
52	7477249CB1	4245	2833-3018, 1869- 2121, 3707-4245, 1-252, 982-1239, 289-357	4669722H1 (SINTNOT24)	1801	2069
				71660072V1	2404	3156
				71657569V1	3106	3854
				7633968J1 (SINTDIE01)	2579	3175
				6440145F8 (BRAENOT02)	938	1087
				71664080V1	3228	3891
				GBI.g8567478.edit	1	2547
				71660176V1	3773	4245
				71662066V1	1802	2475
				2605539F6 (LUNGUT07)	433	939
				71659261V1	1690	2437
				3825558H1 (BRAIHCT02)	1179	1270

WO 02/04520

PCT/US01/21448

Table 4 (cont.)

Seq ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
52				7765571H1 (URRTUE01)	1	693
53	7477720CB1	2124	1-936, 1200- 1488, 1982-2124, 1562-1745	FL7477720-95836195- 9205709	1	2124
54	7477852CB1	2195	1-418, 1899-2195	GB1.98748866.ed1c	1	2195
55	1471717CB1	2055	206-768, 881- 931, 1155-1323	704464956V1 72271206V1	492	297
56	3874406CB1	4727	1-1299, 1576- 1632, 2550-3619, 2014-2192	717938633V1 550522105V1 717988347V1 3620 3575 1380 2475 482	1571	744
57	459654CB1	3852	1-335, 2014-3231	55068154H1 3133035F6 (SMCNOT01) 55058329H1 55068182V1 2048 3593 1778 3348 1666 660 349	223	1718
				6772024J1 (BRANOR01) 55064208J1	1	623

Table 4 (cont.)

Seq ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
57				6617183H2 (BRAXTR14) 6195941H1 (PITUNON01) 71909238V1 2216896F6 (SINTFE03) 71042073V1 7431853H1 GNN:94375937_004-ed 1c	2823	3458
58	5047435CB1	1917	1-238, 1162-1474	6426880H1 (LUNGON07) 6781142H1 (OVARDIR01) 2645767H1 (OVARNOT09) 71704421V1 7726210H1 (THYRDIE01) 7721710J2 (THYRDIE01) 6340173F8 (BNANDIN01) 71704256V1 3025 2408	814	3093
59	7475603CB1	6791	1-3283, 5952- 6101, 3793-4761	71762287V1 (THYMN0E02) 7760618J1 (LIVRFE04) 7462584H1 5817 1 1251 4313	128	4879

This Page is inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked.

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images problems checked, please do not report the problems to the IFW Image Problem Mailbox**

**THIS PAGE BLANK (USPTO)**